

THE KOHN-LAPLACE EQUATION ON ABSTRACT CR MANIFOLDS: LOCAL REGULARITY

TRAN VU KHANH

ABSTRACT. The purpose of this paper is to establish local regularity of the solution operator to the Kohn-Laplace equation, called the complex Green operator, on abstract CR manifolds of hypersurface type. For a cut-off function σ , we introduce the σ -superlogarithmic property, a potential theoretical condition on CR manifolds. We prove that if the given datum is smooth on an open set containing the support of σ then the solution is smooth on the interior of $\{x \in M : \sigma(x) = 1\}$. Furthermore, we also study the smoothness of the integral kernel of the complex Green operator.

In memory of my teacher Beppe Zampieri

CONTENTS

1. Introduction	2
1.1. Introduction and motivation	2
1.2. The main theorems	5
Acknowledgements	10
2. Technical preliminaries	11
2.1. The Kohn-Morrey-Hörmander inequality on CR manifolds	11
2.2. Microlocalization	13
2.3. A special norm between $\ \zeta_0\ _{H^s}$ and $\ \zeta_1\ _{H^s}$	16
3. A-priori estimates	17
3.1. A-priori estimates for the system $(\bar{\partial}_b, \bar{\partial}_b^*)$	18
3.2. A-priori estimates for $\square_b$	24
3.3. A-priori estimates for $\square_b^\delta$	27
4. Proof of the main theorems	29
4.1. Local regularity of the complex Green operator	29
4.2. Smoothness of the complex Green kernel	36
5. The σ -superlogarithmic property on hypersurface models	38
Appendix	45
References	48

Research was supported by the Australian Research Council DE160100173. This work was done in part while the author was a visiting member at the Vietnam Institute for Advanced Study in Mathematics (VIASM). He would like to thank the institution for its hospitality and support.

1. INTRODUCTION

1.1. Introduction and motivation. Let M be an abstract $(2n + 1)$ -dimensional CR manifold of hypersurface type (or CR manifold for short) equipped with a CR structure $T^{1,0}M$. Let θ be a contact form of M which is a nonvanishing purely imaginary 1-form that annihilates $T^{1,0}M \oplus T^{0,1}M$. Then the Levi form on M with respect to θ is the Hermitian form on $T^{1,0}M$ given by $d\theta(L \wedge \bar{L}')$ where $L, L' \in T^{1,0}M$. We say that M is pseudoconvex if the Levi form $d\theta(L \wedge \bar{L})$ is nonnegative for any $L \in T^{1,0}M$. For a C^2 function λ on M , the alternating $(1, 1)$ -form $\mathcal{L}_\lambda := \frac{1}{2}(\partial_b \bar{\partial}_b - \bar{\partial}_b \partial_b)\lambda$ on $T^{1,0}M \times T^{0,1}M$ is called the Levi form of λ .

For $0 \leq q \leq n$, the bundle of $(0, q)$ -forms of smooth coefficients on M is denoted by $C_{0,q}^\infty(M)$. The induced L^2 -inner product and norm on $C_{0,q}^\infty(M)$ is defined by

$$(u, v)_{L^2} = \int_M \langle u, v \rangle dV, \quad \|u\|_{L^2}^2 = (u, u)_{L^2}.$$

where dV is the element of volume on M and $\langle \cdot, \cdot \rangle$ is a Hermitian metric on $T^{1,0}M \oplus T^{0,1}M$. In general, $\|\cdot\|_{H^s}$ denotes the Sobolev H^s -norm that $H^0 := L^2$. We also define $L_{0,q}^2(M)$ and $H_{0,q}^s(M)$ to be the Hilbert space obtained by completing $C_{0,q}^\infty(M)$ under the L^2 and H^s norms, respectively. The tangential Cauchy-Riemann operator $\bar{\partial}_b : C_{0,q}^\infty(M) \rightarrow C_{0,q+1}^\infty(M)$ is the natural restriction of the de-Rham exterior complex of derivatives d to $C_{0,q+1}^\infty(M)$. We denote by $\bar{\partial}_b^* : C_{0,q+1}^\infty(M) \rightarrow C_{0,q}^\infty(M)$ the L^2 -adjoint of $\bar{\partial}_b$ and define the Kohn-Laplacian by

$$\square_b := \bar{\partial}_b \bar{\partial}_b^* + \bar{\partial}_b^* \bar{\partial}_b : C_{0,q}^\infty(M) \rightarrow C_{0,q}^\infty(M).$$

We can extend $\bar{\partial}_b$, $\bar{\partial}_b^*$ and $\square_b$ to unbounded operators in L^2 -spaces with domains consisting of forms in L^2 -spaces where the results (computed in the sense of distributions) are actually in L^2 -spaces. Let $\mathcal{H}_{0,q}(M) = \{u \in L_{0,q}^2(M) : \square_b u = 0\}$ be the space of harmonic $(0, q)$ -forms, that coincides with $\ker(\bar{\partial}_b) \cap \ker(\bar{\partial}_b^*)$.

Given $\varphi \in L_{0,q}^2(M)$ with $\varphi \perp \mathcal{H}_{0,q}(M)$, to solve the Kohn-Laplace equation with datum φ consists in finding a $(0, q)$ -form $u \in \text{Dom}(\square_b)$ such that

$$\square_b u = \varphi. \tag{1.1}$$

Observe that if a solution of (1.1) exists then there is a unique solution u of (1.1) such that $u \perp \mathcal{H}_{0,q}(M)$, we denote the operator that maps $\varphi \mapsto u$ by G_q . The operator G_q is called the complex Green operator acting $(0, q)$ -forms. We extend the operator G_q to a linear operator by setting it to be equal to zero on $\mathcal{H}_{0,q}(M)$. If there exists $c > 0$ so that for any $\varphi \in L_{0,q}^2(M) \cap \mathcal{H}_{0,q}^\perp(M)$, $\|u\|_{L^2} \leq c\|\varphi\|_{L^2}$ then G_q is bounded and self adjoint in $L_{0,q}^2(M)$. Furthermore, if G_q exists, then $\bar{\partial}_b^* G_q$ and $G_q \bar{\partial}_b^*$ are the canonical solution operators to the $\bar{\partial}_b$ -equation on $(0, q)$ and $(0, q + 1)$ -forms, respectively; $\bar{\partial}_b G_q$ and $G_q \bar{\partial}_b$ are the canonical solution operators to the $\bar{\partial}_b^*$ -equation on $(0, q)$ and $(0, q - 1)$ -forms, respectively; $I - \bar{\partial}_b^* \bar{\partial}_b G_q$ and $I - \bar{\partial}_b^* G_q \bar{\partial}_b$ are the Szegő projections—the projections onto

$\ker(\bar{\partial}_b)$; and $I - \bar{\partial}_b \bar{\partial}_b^* G_q$ and $I - \bar{\partial}_b G_q \bar{\partial}_b^*$ are the anti Szegő projections— the projections onto $\ker(\bar{\partial}_b^*)$. The fundamental questions regarding the Kohn-Laplace equation can be asked, at the level of understanding the regularity of these operators, as follows:

- (1) the L^2 boundedness;
- (2) the global regularity in C^∞ and H^s spaces;
- (3) the local regularity in C^∞ and H^s spaces;
- (4) the smoothness of the integral kernels.

A great deal of work has been done relatively to Question 1 (cf. [Bar12, HR11, HR15, Koh86, KN06, Nic06, Sha85]) and Question 2 (cf. [RS08, Rai10, KPZ12, SZ15, Str12, HPR15]) on some classes of CR manifolds. In the parallel with this work in [KR], with Raich we give a positive answer to the first two questions on abstract CR manifolds equipped with “good” CR-plurisubharmonic functions, which is a potential theoretical condition that we call a P -property. In particular, we prove that if M is a compact, pseudoconvex-oriented, $(2n + 1)$ -dimensional CR manifold equipped with a strictly CR-plurisubharmonic function on $(0, q_0)$ -forms, i.e., there exist a global function λ and constant $\alpha > 0$ such that

$$\langle (\mathcal{L}_\lambda + d\theta) \lrcorner u, u \rangle \geq \alpha |u|^2 \quad \text{holds on } M \text{ for all smooth } (0, q)\text{-forms } u,$$

then the operators G_q , $\bar{\partial}_b^* G_q$, $G_q \bar{\partial}_b^*$, $\bar{\partial}_b G_q$, $G_q \bar{\partial}_b$, $I - \bar{\partial}_b^* \bar{\partial}_b G_q$, $I - \bar{\partial}_b^* G_q \bar{\partial}_b$, $I - \bar{\partial}_b \bar{\partial}_b^* G_q$, $I - \bar{\partial}_b G_q \bar{\partial}_b^*$, $\bar{\partial}_b G_q^2 \bar{\partial}_b^*$ and $\bar{\partial}_b^* G_q^2 \bar{\partial}_b$ are L^2 -bounded for all degrees $q_0 \leq q \leq n - q_0$. Here $\lrcorner$ is the contraction operator defined in Section 2 below. If the manifold also admits a covering that satisfies “a weak compactness property” on $(0, q_0)$ -forms, then these operators are both C^∞ -globally regular and exactly regular in H^s -spaces [KR]. Our “weak compactness” property acting $(0, q_0)$ -forms on a covering $\{U_\eta\}_\eta$ of M is defined as follows: for any U_η and any $t \geq 1$ there exist a vector field T_t transversal $T^{1,0}M \oplus T^{0,1}M$, a uniformly bounded, smooth function λ_t^η , and a function $f(t) \nearrow \infty$ as $t \nearrow \infty$ such that

$$\begin{cases} \langle (\mathcal{L}_{\lambda_t^\eta} + td\theta) \lrcorner u, u \rangle \geq f(t) |\{Lie\}_{T_t}(\theta)|^2 |u|^2, \\ 0 < c_1 \leq \theta(T_t) \leq c_2 \quad \text{uniformly in } t, \end{cases} \quad (1.2)$$

hold on U_η for all smooth $(0, q_0)$ -forms u .

The purpose of this paper is to answer Question 3 and 4. The study of the local C^∞ regularity and the smoothness of the distribution kernels of G_q and the relative operators are preliminary steps for the derivation of L^p and Hölder estimates for these operators (cf. [Chr88, Koe02, FK88b, FK88a, NRSW89, NS06, FS74, CNS92, Mac88]). For the question on local regularity, we first recall an operator is called locally regular on a given open set U if it preserve C^∞ on U . The local regularity of G_q is equivalent to the local hypoellipticity of $\square_b$ in the sense that if the restriction of $\square_b u$ to U is in $C_{0,q}^\infty(U)$, then the restriction of u to U is also in $C_{0,q}^\infty(U)$.

It is well known in the general theory of partial differential equations that a subelliptic estimate implies local hypoellipticity [KN65]. For the $\bar{\partial}$ -Neumann problem, Kohn [Koh79]

gave a sufficient condition for subellipticity over pseudoconvex domains in whose boundaries are real analytic and of finite type by introducing a sequence of ideals of subelliptic multipliers. For example, for the hypersurface in $\mathbb{C}^{n+1}$ defined by

$$\text{Im } z_{n+1} = \sum_{k=1}^m |h_k(z_1, \dots, z_n)|^2, \quad (1.3)$$

where h_k 's are holomorphic functions and whose common zeroes is only the origin of $\mathbb{C}^n$, the origin is a point of finite type. Hence, by [Koh79, Theorem 1.19], a subelliptic estimate for $\square$ holds (see [Koh79] for the analysis of this example). In [Cat83, Cat87], Catlin proved, regardless whether the boundary is real analytic or not, the equivalence of the finite type condition and the subelliptic estimate by establishing that each of these two conditions is equivalent to a third technical condition called the *potential-theoretical condition* on domains. For a CR manifold which is the boundary of a bounded pseudoconvex domain in $\mathbb{C}^{n+1}$, the subelliptic estimate for $\square_b$ and $\square$ are equivalent [Koh02]. Consequently, $\square_b$ is locally hypoelliptic on the set of points of finite type. An amazing result by Kohn [Koh02] showed that superlogarithmic estimate for $\square_b$ still implies local hypoellipticity. It should be noted that superlogarithmicity is a very weak estimate compared with subellipticity and that there are many classes of domains of infinite type (flat boundary) for which the superlogarithmic estimate holds (cf. [KZ10]). Superlogarithmicity is also a necessary condition for local hypoellipticity of $\square_b$ on some hypersurfaces. For example, Christ [Chr02] considered the model in $\mathbb{C}^2$ defined by

$$\text{Im } z_2 = \exp \left(-\frac{1}{|\text{Re } z_1|^\alpha} \right), \quad (1.4)$$

for some $\alpha > 0$, and showed that in a neighborhood of the origin, $\square_b$ is locally hypoelliptic if and only if superlogarithmic estimate holds, i.e., $\alpha < 1$ (see also [KZ10, BKZ12]). The problem is to find conditions under which $\square_b$ is locally hypoelliptic but superlogarithmicity fails. It is a recent discovery [Koh00, Chr01, BPZ15] that there exist many classes of hypersurfaces that superlogarithmicity might not hold but $\square_b$ is still locally hypoelliptic. For example, $\square_b$ is locally hypoelliptic on hypersurface models in $\mathbb{C}^{n+1}$ either defined by

$$\text{Im } z_{n+1} = \exp \left(-\frac{1}{(\sum_{k=1}^m |h_k(z_1, \dots, z_n)|^2)^\beta} \right), \quad (1.5)$$

where h_k 's are holomorphic functions whose common zeroes is only at the origin of $\mathbb{C}^n$, and $\beta > 0$ (see [Koh00, Chr01]); or

$$\text{Im } z_{n+1} = \sum_{j=1}^n |\text{Re } z_j|^{2m_j} \exp \left(-\frac{1}{|z_j|^{\beta_j}} \right), \quad (1.6)$$

where $m_j \in \mathbb{N}$ and $\beta_j > 0$, $j = 1, \dots, n$ (see [BPZ15]). The problem of local hypoellipticity of $\square_b$, raised by Zampieri, remains open for the model, defined by

$$\operatorname{Im} z_{n+1} = \sum_{j=1}^n \exp \left(-\frac{1}{|\operatorname{Re} z_j|^{\alpha_j}} \right) \exp \left(-\frac{1}{|z_j|^{\beta_j}} \right) \quad (1.7)$$

where $0 < \alpha_j < 1$ and $\beta_j > 0$, $j = 1, \dots, n$.

The problem raised by Zampieri has motivated my work on this paper. The main goal of this paper consists in establishing the local hypoellipticity of $\square_b$ and hence the regularity of G_q and related operators. The technique is to establish a sufficient potential theoretic condition on abstract CR manifolds. Surprisingly, not only this shows local hypoellipticity of $\square_b$ on the hypersurface defined by (1.7), but also we discover a new class of hypersurfaces in $\mathbb{C}^{n+1}$ in which superlogarithmic estimate for $\square_b$ might not hold but $\square_b$ is locally hypoelliptic. For example, a generalization of Kohn's example in [Koh79]:

$$\operatorname{Im} z_{n+1} = \sum_{j=1}^m H_j(|h_j(z_1, \dots, z_n)|), \quad (1.8)$$

where $H_j : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ are increasing convex functions with $H_j(0) = 0$ and h_j 's are holomorphic functions and have the common zero at only the origin of $\mathbb{C}^n$; or a generalization of Christ's example in [Chr02]:

$$\operatorname{Im} z_2 = \exp \left(-\frac{1}{|\operatorname{Re}(z_1^m)|^\alpha} \right) \quad (1.9)$$

where $m \in \mathbb{N}$ and $0 < \alpha < 1$. In particular, all mentioned examples (1.3)-(1.9) can be unified to a class of hypersurfaces in $\mathbb{C}^{n+1}$ having the form

$$\operatorname{Im} z_{n+1} = \sum_{k=1} H_k \left(\sum_j |h_{kj}|^2 \right) F_k \left(\sum_j |\operatorname{Re} h_{kj}|^2 \right) \quad (1.10)$$

where $H_k, F_k : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ are increasing and convex functions such that $H_k(0)F_k(0) = 0$; and the h_{kj} 's are holomorphic functions in $\mathbb{C}^n$ with an isolated zero at the origin. The crucial condition for local hypoellipticity of $\square_b$ is only that $\lim_{\delta \rightarrow 0} \delta \ln(F_k(\delta^2)) = 0$ and no matter how small the value of H_k 's are.

1.2. The main theorems. Throughout of this paper, we consider M to be a $(2n+1)$ -dimensional, pseudoconvex CR manifold of hypersurface type with a fixed purely imaginary contact form θ and its dual vector field T ; U to be an open set of M on which a smooth basis of $\mathbb{C}TM$ exists. We do not require M to be compact, however, we assume that $\bar{\partial}_b$ has closed range in L^2 for all degree of forms (see [KR] for discussion on this condition). Let σ be a cutoff function, we denote by V_0^σ the interior of the set $\{x \in M : \sigma(x) = 1\}$. If ζ is another cutoff function, then we use notation $\sigma \prec \zeta$ if $\zeta = 1$ on the support of σ . For two functions $f, g : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, we write $f \gg g$ if $\lim_{t \rightarrow \infty} f(t)/g(t) = \infty$.

Now we introduce a potential theoretic condition that is a sufficient condition for local hypoellipticity of $\square_b$.

Definition 1.1. Let σ be a cutoff function in U . We say that the σ -superlogarithmic property on $(0, q_0)$ -forms holds if there exist a pair of rate functions $f, f_0 : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with $f, f_0 \gg 1$ and $f(t), f_0(t) \leq t^{1/2}$ for $t \geq 1$, and a family of smooth real-valued function $\{\lambda_t\}_{t \geq 1}$ defined on U such that

$$\begin{cases} \langle (\mathcal{L}_{\lambda_t} + td\theta) \lrcorner u, u \rangle \geq f_0^2(t) (\ln t |\langle \mathcal{L}_\sigma \lrcorner u, u \rangle| + \ln^2 t |\partial_b \sigma \lrcorner u|^2) + f^2(t) |u|^2, \\ |T^m(\lambda_t)| \leq c_m t^{\frac{m}{2}}, \text{ for all } m \in \mathbb{N}, \end{cases} \quad (1.11)$$

holds at any $x \in U$ for all smooth $(0, q_0)$ -forms u . The functions f, f_0 are called the independent and dependent rate, respectively.

This definition is inspired by the P -property defined by Catlin in [Cat84, Cat87, KZ10] and Property $(CR-P_q)$ defined by Raich in [Rai10, KPZ12, Str12]. The σ -superlogarithmic property is a combination of the potential theoretical conditions $(f-Id)$, $(f_0 \ln -\partial_b \sigma \wedge \bar{\partial}_b \sigma)$ and $(f_0 \sqrt{\ln} - \mathcal{L}_\sigma)$ -properties where the $(f-\mathcal{M})$ -property is defined by the first line of (1.11) replaced by

$$\langle (\mathcal{L}_{\lambda_t} + td\theta) \lrcorner u, u \rangle \geq f^2(t) |\langle \mathcal{M} \lrcorner u, u \rangle|. \quad (1.12)$$

Here, $\mathcal{M}$ is a $(1, 1)$ form and is called a multiplier of the f -property. It is automatic that $(t^{1/2} - d\theta)$ -property holds on any CR manifold. A byproduct from the proof of the main result in §3.1, we have a general estimate for $\square_b$ by the $(f-\mathcal{M})$ -property.

Theorem 1.2. Let M be a $(2n+1)$ -dimensional, pseudoconvex CR manifold and $x_o \in M$, let $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, and $\mathcal{M}$ be a nonnegative $(1, 1)$ -form. Suppose $(f-\mathcal{M})$ -property on $(0, q_0)$ -forms holds at in a neighborhood U of x_o . Then, the estimate

$$(\mathcal{M} \lrcorner f(\Lambda) u^+, f(\Lambda) u^+)_{L^2} \leq c (\|\bar{\partial}_b u^+\|_{L^2}^2 + \|\bar{\partial}_b^* u^+\|_{L^2}^2 + \|u\|_{L^2}^2) \quad (1.13)$$

holds for all $u \in C_{0,q}^\infty(M)$ supported in U with $q \geq q_0$; and dually, the estimate

$$\begin{aligned} \|\sqrt{\text{Tr}(\mathcal{M})} \times f(\Lambda) u^-, f(\Lambda) u^-\|_{L^2}^2 - (\mathcal{M} \lrcorner f(\Lambda) u^-, f(\Lambda) u^-)_{L^2} \\ \leq c (\|\bar{\partial}_b u^-\|_{L^2}^2 + \|\bar{\partial}_b^* u^-\|_{L^2}^2 + \|u\|_{L^2}^2) \end{aligned} \quad (1.14)$$

holds for all $u \in C_{0,q}^\infty(M)$ supported in U with $q \leq n - q_0$. Here $u^\pm$ are the $\pm$ -Kohn's microlocalizations of u and $f(\Lambda)$ is the pseudodifferential operator of symbol $f(\sqrt{1 + |\xi|^2})$.

We remark that if $\mathcal{M}$ is a positive $(1, 1)$ -form, then (1.13), (1.14) and the elliptic estimate for the 0-Kohn's microlocalization u^0 imply the full estimate

$$\|f(\Lambda) u\|_{L^2}^2 \leq c (\|\bar{\partial}_b u\|_{L^2}^2 + \|\bar{\partial}_b^* u\|_{L^2}^2 + \|u\|_{L^2}^2) \quad (1.15)$$

for all $u \in C_{0,q}^\infty(M)$ supported in U with $q_0 \leq q \leq n - q_0$. In this situation, if $f(t) = t^\epsilon$, (1.15) becomes a subelliptic estimate; if $f \gg \ln$, (1.15) implies the superlogarithmic estimate; and if $f \gg 1$, (1.15) implies the (local) compactness estimate. These estimates are

well-known in theory of hypoellipticity of $\square_b$. The compactness estimate for $\square_b$ obtained by potential-theoretical property has been proved in [Rai10, Str12, KPZ12]. However, this is the first time that estimates for $\square_b$ by potential-theoretical properties have shown a “gain” in derivatives. Coming back to Definition (1.1), it is obvious that if the independent rate $f \gg \ln$ then the σ -superlogarithmic property holds for any cutoff function σ in U . In fact, in this case the superlogarithmic estimate for $\square_b$ holds by Theorem 1.2 and hence $\square_b$ is local hypoellipticity by [Koh02, Theorem 1.5].

The σ -superlogarithmic property implies the following results concerning hypoellipticity of $\square_b$ and regularity of G_q and its related operators on CR manifolds with dimension at least five.

Theorem 1.3. *Let M be a pseudoconvex CR manifold of dimension $(2n+1)$ with $n \geq 2$ such that $\bar{\partial}_b$ has closed range in L^2 spaces for all degrees of forms. Suppose that the σ -superlogarithmic property on $(0, q_0)$ -forms holds. Then, for any $q_0 \leq q \leq n - q_0$, if $u \in L^2_{0,q}(M)$ such that $\square_b u \in L^2_{0,q}(M) \cap C^\infty_{0,q}(V_1^\sigma)$ then $u \in C^\infty_{0,q}(V_0^\sigma)$, where V_0^σ the interior of $\{x \in M : \sigma(x) = 1\}$ and V_1^σ be an open set containing $\text{supp}(\sigma)$.*

Furthermore, for any $s \geq 0$, and $\zeta_0 \prec \sigma \prec \zeta_1$, the following holds.

(i) *If $\varphi \in L^2_{0,q}(M) \cap H^s_{0,q}(\text{supp}(\zeta_1))$, then*

$$\begin{aligned} & \|f^2(\Lambda)\zeta_0 G_q \varphi\|_{H^s} + \|f(\Lambda)\zeta_0 \bar{\partial}_b G_q \varphi\|_{H^s} + \|f(\Lambda)\zeta_0 \bar{\partial}_b^* G_q \varphi\|_{H^s} \\ & + \|\zeta_0(I - \bar{\partial}_b \bar{\partial}_b^* G_q) \varphi\|_{H^s} + \|\zeta_0(I - \bar{\partial}_b^* \bar{\partial}_b G_q) \varphi\|_{H^s} \leq c_{s,\zeta_0,\zeta_1} (\|\zeta_1 \varphi\|_{H^s} + \|\varphi\|_{L^2}). \end{aligned} \quad (1.16)$$

(ii) *If $\varphi \in L^2_{q-1}(M) \cap H^s_{0,q-1}(\text{supp}(\zeta_1))$, then*

$$\begin{aligned} & \|f^2(\Lambda)\zeta_0 \bar{\partial}_b^* G_q^2 \bar{\partial}_b \varphi\|_{H^s} + \|f(\Lambda)\zeta_0 G_q \bar{\partial}_b \varphi\|_{H^s} + \|\zeta_0(I - \bar{\partial}_b^* G_q \bar{\partial}_b) \varphi\|_{H^s} \\ & \leq c_{s,\zeta_0,\zeta_1} (\|\zeta_1 \varphi\|_{H^s} + \|\varphi\|_{L^2}). \end{aligned} \quad (1.17)$$

(iii) *If $\varphi \in L^2_{0,q+1}(M) \cap H^s_{0,q+1}(\text{supp}(\zeta_1))$,*

$$\begin{aligned} & \|f^2(\Lambda)\zeta_0 \bar{\partial}_b G_q^2 \bar{\partial}_b^* \varphi\|_{H^s} + \|f(\Lambda)\zeta_0 G_q \bar{\partial}_b^* \varphi\|_{H^s} + \|\zeta_0(I - \bar{\partial}_b G_q \bar{\partial}_b^*) \varphi\|_{H^s} \\ & \leq c_{s,\zeta_0,\zeta_1} (\|\zeta_1 \varphi\|_{H^s} + \|\varphi\|_{L^2}). \end{aligned} \quad (1.18)$$

Here, f is the independent rate of the σ -superlogarithmic property.

The proof of Theorem 1.3 is given in §4.1. The crucial idea in the proof is to combine the elliptic regularization method and a-priori estimates in the intermediate norm $\|\cdot\|_{f-\mathcal{A}^{s\sigma}_\zeta}$ between two classical norms $\|f(\Lambda)\zeta_0 \cdot\|_{H^s}$ and $\|f(\Lambda)\zeta_1 \cdot\|_{H^s}$. In particular, we will prove that

$$\|u\|_{f-\mathcal{A}^{s\sigma}_\zeta} \leq c_{s,\sigma,\zeta} \left(\|\bar{\partial}_b u\|_{\mathcal{A}^{s\sigma}_\zeta} + \|\bar{\partial}_b^* u\|_{\mathcal{A}^{s\sigma}_\zeta} + \|u\|_{L^2} \right),$$

and

$$\begin{aligned} & \|u\|_{f^2-\mathcal{A}_\zeta^\sigma} + \|\bar{\partial}_b u\|_{f-\mathcal{A}_\zeta^\sigma} + \|\bar{\partial}_b^* u\|_{f-\mathcal{A}_\zeta^\sigma} + \|\bar{\partial}_b \bar{\partial}_b^* u\|_{\mathcal{A}_\zeta^\sigma} + \|\bar{\partial}_b^* \bar{\partial}_b u\|_{\mathcal{A}_\zeta^\sigma} \\ & \leq c_{s,\sigma,\zeta} \left(\|\square_b^\delta u\|_{\mathcal{A}_\zeta^\sigma} + \|u\|_{L^2} \right) \end{aligned}$$

hold for all $u \in L_{0,q}^2(M) \cap C_{0,q}^\infty(\text{supp}(\zeta))$. Here $\square_b^\delta$ is an elliptic perturbation of $\square_b$. It is should be emphasized that with the stronger hypotheses in [Chr01, Chr02, Koh02, Koh00, BPZ15] the difficulty of regularization methods can be avoided.

Remark 1.4. For the top degrees, it is known that $G_0 := \bar{\partial}_b^* G_1^2 \bar{\partial}_b$ and $G_n = \bar{\partial}_b G_{n-1}^2 \bar{\partial}_b^*$. Thus, if the σ -superlogarithmic property holds on $(0,1)$ -forms then by Theorem 1.3 (ii) and (iii) we get

$$\|f^2(\Lambda) \zeta_0 G_0 \varphi\|_{H^s} + \|f(\Lambda) \zeta_0 \bar{\partial}_b G_0 \varphi\|_{H^s} \leq c_{s,\zeta_0,\zeta_1} (\|\zeta_1 \varphi\|_{H^s} + \|\varphi\|_{L^2}). \quad (1.19)$$

holds for all $\varphi \in L^2(M) \cap H^s(\text{supp}(\zeta_1))$, and

$$\|f^2(\Lambda) \zeta_0 G_n \varphi\|_{H^s} + \|f(\Lambda) \zeta_0 \bar{\partial}_b^* G_n \varphi\|_{H^s} \leq c_{s,\zeta_0,\zeta_1} (\|\zeta_1 \varphi\|_{H^s} + \|\varphi\|_{L^2}). \quad (1.20)$$

holds for all $\varphi \in L_{0,n}^2(M) \cap H_{0,n}^s(\text{supp}(\zeta_1))$. However, this works only for CR manifolds of dimension at least five, i.e, $n \geq 2$. For the case $n = 1$, we add an extra assumption that G_0 is C^∞ globally regular. Here is our result for this case.

Theorem 1.5. *Let M be a 3-dimensional pseudoconvex CR manifold such that $\bar{\partial}_b$ has closed range on functions and G_0 is globally regular. Suppose that the σ -superlogarithmic property with dependent rate f holds for $(0,1)$ -forms. Then, for $q = 0, 1$, if $u \in L_{0,q}^2(M)$ such that $\square_b u \in L_{0,q}^2(M) \cap C_{0,q}^\infty(V_1^\sigma)$ then $u \in C_{0,q}^\infty(V_0^\sigma)$. Furthermore, (1.19) and (1.20) hold.*

Let $\mathcal{G}_q(\hat{x}, x)$ be the integral kernel of the complex Green operator G_q . This means G_q has the integral representation

$$G_q \varphi(x) = \int_M \langle \varphi(\hat{x}), \mathcal{G}_q(\hat{x}, x) \rangle dV(\hat{x}) = (\varphi, \mathcal{G}_q(\cdot, x))$$

provided the integral exists; this imposes regularity conditions on φ and $\mathcal{G}_q$. Thus, G is a double differential form of degree $(0, q; q, 0)$ on the product manifold $M \times M$. The second goal of this paper is to study the smoothness of $\mathcal{G}_q$. Using the idea of Kerzman [Ker72] for proving smoothness of the Bergman kernel of strongly pseudoconvex domains in $\mathbb{C}^{n+1}$, the smoothness of the Szegő kernels and the kernel of the complex Green operator have been obtained on some cases of pseudoconvex hypersurfaces in $\mathbb{C}^{n+1}$ of finite type (see [MS97, Chr88, NRSW89, PS77, Boa87, NS06, Koe02, FKM90]). However, when M is of infinite type, the only result available is that of Halfpap, Nagel and Wainger [HNW10] who classified subsets of $M \times M$ in which the Szegő kernel is smooth by comparing α with 1 for the case that M is defined by $\text{Im } z_2 = \exp\left(-\frac{1}{|\text{Re } z_1|^\alpha}\right)$. Halfpap et. al. use a harmonic analysis approach.

It is obvious that the smoothness of $\mathcal{G}_q$ implies the smoothness of the Szegő kernels of $I - \bar{\partial}_b^* \bar{\partial}_b G_q$ and $I - \bar{\partial}_b^* G_q \bar{\partial}_b$ and also the kernels of all related operators such as $\bar{\partial}_b G_q$, $G_q \bar{\partial}_b$, $\bar{\partial}_b^* G_q$, $G_q \bar{\partial}_b^*$, $I - \bar{\partial}_b \bar{\partial}_b^* G_q$, and $I - \bar{\partial}_b G_q \bar{\partial}_b^*$. So we only need to state the result on the smoothness of G_q .

Theorem 1.6. *Let M be a pseudoconvex, CR manifold of dimension $(2n+1)$ such that $\bar{\partial}_b$ has closed range in L^2 spaces for all degrees of forms. Suppose that the σ -superlogarithmic property on $(0, q_0)$ forms holds .*

- (i) *If $\hat{\sigma}$ is another cutoff function such that $\text{supp}(\sigma) \cap \text{supp}(\hat{\sigma}) = \emptyset$ and the $\hat{\sigma}$ - superlogarithmic property on $(0, q_0)$ -forms also holds. Then*

$$\mathcal{G}_q \in C_{0,q;0}^\infty \left((V_0^\sigma \times V_0^{\hat{\sigma}}) \cup (V_0^{\hat{\sigma}} \times V_0^\sigma) \right) \quad \text{for all } q_0 \leq q \leq n - q_0.$$

If $q_0 = 1$ and $n \geq 2$, or $q_0 = n = 1$ with the extra assumption that G_0 is globally regular, then this conclusion also holds for $q = 0$ and $q = n$.

- (ii) *If G_q is globally regular for some $q_0 \leq q \leq n - q_0$ then*

$$\mathcal{G}_q \in C_{0,q;q,0}^\infty \left(\left(V_0^\sigma \times (M \setminus \overline{\text{supp}(\sigma)}) \right) \cup \left((M \setminus \overline{\text{supp}(\sigma)}) \times V_0^\sigma \right) \right).$$

If $q_0 = q = 1$ (resp. $q_0 = 1, q = n$), this conclusion also holds for G_0 and (resp. G_n).

We prove this theorem in §4.2. The idea of the proof is based on the estimates of regularity for $G_q \varphi_{\hat{x}_0}$ and $G_q^\delta \varphi_{\hat{x}_0}$ (where G_q^δ is the inverse of $\square_b^\delta -$ a elliptic perturbation of $\square_b$) where $\varphi_{\hat{x}_0}$ is a Dirac-delta distribution $(0, q)$ -form supported in $\hat{x}_0$.

We notice that the regularity of G_q and the smoothness of $\mathcal{G}_q$ in Theorem 1.3, 1.5 and 1.6 depends on the set $V_0^\sigma = \{x \in M : \sigma(x) = 1\}$ and the support of σ . To give the precise result for local regularity defined above, we define the point of weak superlogarithmicity as follows.

Definition 1.7. The point $x_0 \in M$ is called a weakly superlogarithmic point if given any small neighborhood W of x_0 then there exists a cutoff function σ such that

- (i) $\text{supp}(\sigma) \subset W$;
- (ii) $\sigma = 1$ in an open set containing x_0 ;
- (iii) the σ -superlogarithmic property on $(0, 1)$ -forms holds.

Denote by S the set of all weakly superlogarithmic points. If $S \subset \subset M$ then S is open. The following corollary follows immediately from Theorems 1.3, 1.5 and 1.6.

Corollary 1.8. *Let M be a pseudoconvex, CR manifold of dimension $(2n+1)$ such that $\bar{\partial}_b$ has closed range in L^2 spaces for all degrees of forms (here $n \geq 2$ or in the case $n = 1$ with the additional hypothesis that G_0 is globally regular). Let S be the set of all weakly superlogarithmic points in M . Then, the following properties hold:*

- (i) *For all $0 \leq q \leq n$, G_q is locally regular in S in the sense that for any open set $V \subset S$, if $\varphi \in L_{0,q}^2(M) \cap C_{0,q}^\infty(V)$ then $G_q \varphi \in C_{0,q}^\infty(V)$.*

- (ii) The kernel $\mathcal{G}_q \in C_{0,q;q,0}^\infty(S \times S \setminus \{\text{Diagonal}\})$ for all $0 \leq q \leq n$.
- (iii) If G_q is global regular for some q , then $\mathcal{G}_q \in C_{0,q;q,0}^\infty((S \times M) \cup (M \times S) \setminus \{\text{Diagonal}\})$.

As our motivation, we will show any point of the hypersurface defined by (1.10) is a weakly superlogarithmic point and then $\square_b$ is locally hypoelliptic.

Theorem 1.9. *Let M^{2n+1} be the hypersurface defined by (1.10). If*

$$\lim_{\delta \rightarrow 0} \delta \ln(F_k(\delta^2)) = 0 \quad \text{for all } k,$$

then any point $x \in M$ is a weakly superlogarithmic point. Consequently, $\square_b$'s defined on forms of any degree are local hypoelliptic.

The hypoellipticity of $\square_b$ acting on functions implies the hypoellipticity of the sublaplacian $\Delta_b := \text{Re}(\square_b)$. The study of Δ_b is related to the work on degenerate second-order operators. By the celebrated theorem of Hörmander [Hör67] the finite type condition of $\{X_0, \dots, X_r\}$ is sufficient for hypoellipticity of $\sum_{j=1}^r X_j^2 + X_0 + c$ in $\mathbb{R}^N$ (see also in [FK72], [RS76]). However, the finite type is not a necessary condition for local hypoellipticity (see [Fed71], [Koh98], [BM95], [KS87], [Mor87], [Him95]). For example, Fedii [Fed71] showed that the operator $\frac{\partial^2}{\partial x^2} + b(x)\frac{\partial^2}{\partial t^2}$ in $\mathbb{R}^2$ with $b \in C^\infty$ is hypoelliptic if $b(x) > 0$ when $x \neq 0$; or the example of Kusuoka-Stroock [KS87] illustrates that the operator $\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + a^2(x)\frac{\partial^2}{\partial t^2}$ in $\mathbb{R}^3$ is hypoelliptic if and only if $\lim_{x \rightarrow 0} x \ln a(x) = 0$, where $a \in C^\infty$, $a(x) \neq 0$ when $x \neq 0$. Applying Theorem 1.9 to the model $\text{Im} z_2 = H(|z_1|^2)F(|\text{Re } z_1|^2)$ in $\mathbb{C}^2$, we obtain a generalized version of Fedii and Kusuoka-Stroock's examples. In fact, we can conclude that the operator $\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + a^2(x)b(x^2 + y^2)\frac{\partial^2}{\partial t^2}$ in $\mathbb{R}^3$ is hypoelliptic if $\lim_{x \rightarrow 0} x \ln a(x) = 0$.

The rest of the paper is organized as follows. The establishment of a basic estimate and the development of the microlocal are given in the first and second parts of Section 2. The remainder of the section introduces a special norm between the classical H^s -norms. In Section 3, we establish the a-priori estimates for the system $(\bar{\partial}_b, \bar{\partial}_b^*)$, the Kohn-Laplacian $\square_b$ and the elliptic perturbation $\square_b^\delta$. We prove Theorems 1.3, 1.5 and 1.6 in Section 4. In Section 5, we analyze the hypersurface model 1.10 and prove Theorem 1.9.

Acknowledgements. I am grateful to Joseph J. Kohn for his scientific discoveries. This research is inspired from his work. I wish to express my thanks to Luca Baracco and Stefano Pinton, communication with them led to the research presented here. I also thank Andy Raich for going through the original manuscript and suggesting several improvements and clarifications.

2. TECHNICAL PRELIMINARIES

In this section we prepare some inequalities which are needed for the *a-priori* estimates that prove Theorems 1.3 and 1.5. The key technical tool of our discussion is the Kohn-Morrey-Hörmander inequality on abstract CR manifolds, the Kohn-microlocalizations and a special norm between $\|\zeta_0\|_{H^s}^2$ and $\|\zeta_1\|_{H^s}^2$.

2.1. The Kohn-Morrey-Hörmander inequality on CR manifolds. Let M be a smooth CR manifold of dimension $(2n+1)$ with the CR structure $T^{1,0}M$ and contact form θ . Let U be an open set of M so that we can choose a $C^\infty(U)$ local basis $\{L_1, \dots, L_n, \bar{L}_1, \dots, \bar{L}_n, T\}$ of $\mathbb{C}TM$. Here, $\{L_1, \dots, L_n\}$ is a basis of $T^{1,0}M$, $\{\bar{L}_1, \dots, \bar{L}_n\}$ is the basis of the conjugate of $T^{1,0}M$, denoted by $T^{0,1}M$. The vector field T is dual to θ . Then the coefficients of the Levi form can be written as $c_{ij} = d\theta(L_i \wedge \bar{L}_j)$, and by the Cartan formula it is identified with the T -component of $[L_i, \bar{L}_j]$. For a smooth function λ , the matrix (λ_{ij}) of the Hermitian form $\mathcal{L}_\lambda := \frac{1}{2}(\partial_b \bar{\partial}_b - \bar{\partial}_b \partial_b)\lambda$ is expressed by

$$\lambda_{ij} = \bar{L}_j L_i(\lambda) + \frac{1}{2} c_{ij} T(\lambda) + \sum_k c_{ij}^k L_k(\lambda). \quad (2.1)$$

where c_{ij}^k is the L_k -component of $[L_i, \bar{L}_j]$ (see [KR] for more details).

To express a form in local coordinates, we let $\mathcal{I}_q = \{(j_1, \dots, j_q) \in \mathcal{N}^q : 1 \leq j_1 < \dots < j_q \leq n\}$, and for $J \in \mathcal{I}_q$, $I \in \mathcal{I}_{q-1}$, and $j \in \mathbb{N}$, ϵ_j^{jI} be the sign of the permutation $\{j, I\} \rightarrow J$ if $\{j\} \cup I = J$ as sets, and 0 otherwise. If $u \in C_{0,q}^\infty(M)$, then u is locally expressed as a combination

$$u = \sum_{J \in \mathcal{I}_q} u_J \bar{\omega}_J, \quad (2.2)$$

of basis forms $\bar{\omega}_J = \bar{\omega}_1 \wedge \dots \wedge \bar{\omega}_{j_k}$ for ordered indices $j_1 < \dots < j_k$ with C^∞ -coefficients u_J . If $\lambda = \sum_j \lambda_j \omega_j$ is a $(1, 0)$ form defined in U , the contraction operator $\lrcorner$ acting $(0, q)$ -form u as in (2.2) is defined by

$$\lambda \lrcorner u = \sum_{I \in \mathcal{I}_{q-1}} \left(\sum_{j=1}^n \lambda_j u_{jI} \right) \bar{\omega}_I, \quad (2.3)$$

and consequently

$$\|\lambda \lrcorner u\|^2 = \sum_{I \in \mathcal{I}_{q-1}} \left\| \sum_{j=1}^n \lambda_j u_{jI} \right\|^2, \quad (2.4)$$

If $\lambda = \sum_{ij} \lambda_{ij} \omega_i \wedge \bar{\omega}_j$ is a $(1, 1)$ form in U , the contraction operator $\lrcorner$ acting a $(0, q)$ -form u as in (2.2) is defined by

$$\lambda \lrcorner u = \sum_{I \in \mathcal{I}_{q-1}} \left(\sum_{i,j=1}^n \lambda_{ij} u_{iI} \right) \bar{\omega}_j \wedge \bar{\omega}_I. \quad (2.5)$$

Thus, the Levi forms $d\theta$ and $\mathcal{L}_\lambda$ acting on $(0, q)$ -forms u, v defined in U can be expressed as

$$\langle d\theta \lrcorner u, v \rangle = \sum_{I \in \mathcal{I}_{q-1}} \sum_{i,j=1}^n c_{ij} u_i \bar{v}_{jI} \quad \text{and} \quad \langle \mathcal{L}_\lambda \lrcorner u, v \rangle = \sum_{I \in \mathcal{I}_{q-1}} \sum_{i,j=1}^n \lambda_{ij} u_i \bar{v}_{jI}.$$

We can also express the operator $\bar{\partial}_b : C_{0,q}^\infty(M) \rightarrow C_{0,q+1}^\infty(M)$ and $\bar{\partial}_b^* : C_{0,q}^\infty(M) \rightarrow C_{0,q-1}^\infty(M)$ in the local basis as follows:

$$\bar{\partial}_b u = \sum_{\substack{J \in \mathcal{I}_q \\ K \in \mathcal{I}_{q+1}}} \sum_{k=1}^n \epsilon_K^{kJ} \bar{L}_k u_J \bar{\omega}_K + \sum_{\substack{J \in \mathcal{I}_q \\ K \in \mathcal{I}_{q+1}}} b_{JK} u_J \bar{\omega}_K \quad (2.6)$$

and

$$\bar{\partial}_b^* u = - \sum_{J \in \mathcal{I}_{q-1}} \left(\sum_{j=1}^n L_j u_{jJ} + \sum_{K \in \mathcal{I}_q} a_{JK} u_K \right) \bar{\omega}_J \quad (2.7)$$

where $a_{ijI}^J, a_{JK} \in C^\infty(M)$ and u is as (2.2).

We will use the Kohn-Morrey-Hörmander inequality for CR manifolds with the weighted norm $\|\cdot\|_{L_\lambda^2}$ defined by

$$\|u\|_{L_\lambda^2}^2 = \int_M \langle u, u \rangle e^{-\lambda} dV.$$

Let $\bar{\partial}_{b,\lambda}^*$ be the L_λ^2 -adjoint of $\bar{\partial}_b$. It is easy to see that in the local basis

$$\bar{\partial}_{b,\lambda}^* u = - \sum_{J \in \mathcal{I}_{q-1}} \left(\sum_{j=1}^n L_j^\lambda u_{jJ} + \sum_{K \in \mathcal{I}_q} a_{JK} u_K \right) \bar{\omega}_J \quad (2.8)$$

where $L_j^\lambda \varphi := e^\lambda L_j(e^{-\lambda} \varphi)$ and u as in (2.2). For such u ,

$$\partial_b(\lambda) \lrcorner u = -[\bar{\partial}_b^*, \lambda]u = [\bar{\partial}_b, \lambda]^* u = \sum_{I \in \mathcal{I}_{q-1}} \sum_{j=1}^n L_j(\lambda) u_{jI} \bar{\omega}_I,$$

and hence

$$\bar{\partial}_{b,\lambda}^* u = \bar{\partial}_b^* u - \partial_b \lambda \lrcorner u. \quad (2.9)$$

Now we can state the Kohn-Morrey-Hörmander inequality type for CR manifold which has been proven in [KR].

Proposition 2.1. *Let λ be a real-valued C^2 function defined on U . Then there exists a constant c independent of λ such that*

$$\begin{aligned}
\|\bar{\partial}_b u\|_{L_\lambda^2}^2 + \|\bar{\partial}_{b,\lambda}^* u\|_{L_\lambda^2}^2 + c\|u\|_{L_\lambda^2}^2 &\geq \frac{1}{2} \sum_{j=1}^n \|\bar{L}_j u\|_{L_\lambda^2}^2 \\
&+ (\mathcal{L}_\lambda \lrcorner u, u)_{L_\lambda^2} + \operatorname{Re} \left\{ (d\theta \lrcorner T u, u)_{L_\lambda^2} \right\}
\end{aligned} \tag{2.10}$$

holds for all $u \in C_{0,q}^\infty(M)$ supported in U .

To estimate the the Levi form containing T in the inequality (2.10), we use Kohn's microlocalizations and Gårding's inequality. Gårding's inequality is formulated as follows (see [Nic06]).

Lemma 2.2. *Let R be a first-order pseudodifferential operator such that the symbol $\mathcal{S}(R) \geq 0$ and let λ be a nonnegative $(1,1)$ -form. Then there exists a constant c such that*

$$\operatorname{Re} \{ (\lambda \lrcorner R u, u)_{L^2} \} \geq -c\|u\|_{L^2}^2$$

for any $u \in C_{0,q}^\infty(M)$.

2.2. Microlocalization. We now recall the microlocal framework developed by Kohn [Koh86, Koh02] and present its main consequences for our work.

We choose real coordinates $x = (x', x_{2n+1}) = (x_1, \dots, x_{2n}, x_{2n+1})$ with the origin some point $x_o \in U$ such that $T = -\sqrt{-1} \frac{\partial}{\partial x_{2n+1}}$ and if we set $z_j = x_j + \sqrt{-1} x_{j+n}$ for $j = 1, \dots, n$, then we have $L_j|_{x_o} = \frac{\partial}{\partial z_j}$ for $j = 1, \dots, n$. Let $\xi = (\xi_1, \dots, \xi_{2n+1}) = (\xi', \xi_{2n+1})$ be the dual coordinates to the $(x_1, \dots, x_{2n+1})$'s. Let ψ be a smooth function in the unit sphere $\{|\xi| = 1\}$ with range in $[0, 1]$, such that

$$\psi = 1 \text{ in } \{\xi | \xi_{2n+1} > \frac{1}{3} |\xi'|\} \quad \text{and} \quad \operatorname{supp}(\psi) \subset \subset \{\xi_{2n+1} \geq \frac{1}{4} |\xi'|\}.$$

We extend this function to $\mathbb{R}^{2n+1}$ by homogeneity. Set $\psi(\xi) := \psi(\frac{\xi}{|\xi|})$ if $|\xi| \geq 2$ and $\psi(\xi) = 0$ if $|\xi| \leq 1$. Set $\psi^+(\xi) := \psi(\xi)$; $\psi^-(\xi) := \psi(-\xi)$ and $\psi^0(\xi) := 1 - \psi^+(\xi) - \psi^-(\xi)$. Define $\tilde{\psi}^0$ so that $\tilde{\psi}^0 = 1$ on a neighborhood of $\operatorname{supp} \psi^0 \cup \operatorname{supp}(d\psi^+) \cup \operatorname{supp}(d\psi^-)$ and $\operatorname{supp}(\tilde{\psi}^0) \subset \mathcal{C}^0 := \{\xi_{2n+1} \leq \frac{3}{4} |\xi'|\} \cup \{|\xi| < 2\}$.

Associated to a function ψ is a pseudodifferential operator Ψ with symbol $\mathcal{S}(\Psi) = \psi$ whose action on a function v supported in U is defined by

$$\widehat{\Psi v}(\xi) = \psi(\xi) \hat{v}(\xi),$$

where $\hat{\cdot}$ denotes the Fourier transform. We say that a pseudodifferential operator Ψ is dominated by a pseudodifferential operator $\tilde{\Psi}$ and denote by $\Psi \prec \tilde{\Psi}$ if $\mathcal{S}(\Psi) \prec \mathcal{S}(\tilde{\Psi})$. The operators Ψ^+ , Ψ^- , Ψ^0 and $\tilde{\Psi}^0$ are defined as above with symbols ψ^+ , ψ^- , ψ^0 and $\tilde{\psi}^0$, respectively. By the definition of ψ^+ , ψ^- , ψ^0 and $\tilde{\psi}^0$, it follows that $\Psi^+ + \Psi^- + \Psi^0 = I$, and both operators $[\Psi^\pm, \alpha]$ and Ψ^0 are dominated by $\tilde{\Psi}^0$, where $\alpha \in C^\infty(M)$.

Let $\gamma : \mathbb{R} \rightarrow [0, 1]$ be a cutoff function such that

$$\gamma(t) = \begin{cases} 1 & \text{if } |t| \in [e^2, e^3], \\ 0 & \text{if } |t| \in [0, e] \cup [e^4, +\infty). \end{cases}$$

Set $\gamma_k(\xi) := \gamma(e^{-k}|\xi_{2n+1}|)$ for $k = 1, 2, \dots$, and also let $\gamma_0(\zeta) = 1$ if $|\zeta_{2n+1}| \leq e$ and is zero if $|\zeta_{2n+1}| \geq e^2$. Furthermore, we can choose γ and γ_0 such that $\sum_{k=0}^{\infty} \gamma_k(\xi) = 1$ for any $\xi \in \mathbb{R}^{2n+1}$. For $k = 0, 1, 2, \dots$, denote by Γ_k the pseudodifferential operator with symbol γ_k . Let ζ be a cutoff function in U and set

$$u_{\zeta,k}^+ := \zeta \Gamma_k \Psi^+ \zeta u, \quad \text{and} \quad u_{\zeta,k}^- := \zeta \Gamma_k \Psi^- \zeta u,$$

where u is a distribution function on M .

Remark 2.3. For $u \in H^{-s_o}(M)$, Plancherel's theorem gives us

$$\begin{aligned} \|u_{\zeta,k}^\pm\|_{H^s}^2 &\leq c \int_{\xi \in \mathbb{R}^{2n+1}} (1 + |\xi|^2)^s \left(\gamma(e^{-k}|\xi_{2n+1}|) \psi^\pm(\xi) \right)^2 |\widehat{\zeta u}(\xi)|^2 d\xi \\ &\leq c_{k,s,r} \int_{\xi \in \mathbb{R}^{2n+1}} (1 + |\xi|^2)^{-s_o} |\widehat{\zeta u}(\xi)|^2 d\xi \\ &= c_{k,s,r} \|\zeta u\|_{H^{-r}}^2 \leq c_{k,s,s_o} \|u\|_{H^{-s_o}}^2, \end{aligned}$$

for all $s, r \in \mathbb{R}$ with $r \geq s_o$. Consequently, by the Sobolev lemma, the functions $u_{\zeta,k}^\pm \in C^\infty(M)$ if $u \in H^{-s_o}(M)$. By this remark we will not worry about the regularity of $u_{\zeta,k}^\pm$.

We finish this subsection by an estimate for the Levi form involving T on the positive microlocalizations $u_{\zeta,k}^+$. The estimate on the negative microlocalizations $u_{\zeta,k}^-$ will be obtained by a modified Hodge-* constructed in §3 below.

Proposition 2.4. *Let $\lambda := \lambda_{k,s}$ be a C^∞ real-valued function defined on U such that $|\lambda| \leq c_{0,s}k$ and $|T^m(\lambda)| \leq c_{m,s}e^{\frac{km}{2}}$ for all $m = 1, 2, \dots$. Then, for $u \in H_{0,q}^{-s_o}(M)$, $r \geq s_o$ and $k \geq 1$, we have*

$$\operatorname{Re} (d\theta \lrcorner T u_{\zeta,k}^+, u_{\zeta,k}^+)_{L_\lambda^2} \geq \frac{1}{2} e^k (d\theta \lrcorner u_{\zeta,k}^+, u_{\zeta,k}^+)_{L_\lambda^2} - c \|u_{\zeta,k}^+\|_{L_\lambda^2}^2 - c_{s,r} e^{-k} \|\zeta u\|_{H^{-r}}^2. \quad (2.11)$$

where c and $c_{s,r}$ are independent of k .

Proof. Since T is a purely imaginary operator and λ is a real-valued function, it follows $\operatorname{Re} \left\{ (T(\lambda) d\theta \lrcorner u_{\zeta,k}^+, u_{\zeta,k}^+)_{L_\lambda^2} \right\} = 0$. Hence, we can bring $e^{-\frac{\lambda}{2}}$ inside T and rewrite the LHS of (2.11) as follows

$$\begin{aligned} \operatorname{Re} \left\{ (d\theta \lrcorner T u_{\zeta,k}^+, u_{\zeta,k}^+)_{L_\lambda^2} \right\} &= \operatorname{Re} \left\{ \left(d\theta \lrcorner T e^{-\frac{\lambda}{2}} u_{\zeta,k}^+, e^{-\frac{\lambda}{2}} u_{\zeta,k}^+ \right)_{L^2} \right\} \\ &= \operatorname{Re} \left\{ \left(d\theta \lrcorner (T - e^k) e^{-\frac{\lambda}{2}} u_{\zeta,k}^+, e^{-\frac{\lambda}{2}} u_{\zeta,k}^+ \right)_{L^2} \right\} + e^k (d\theta \lrcorner u_{\zeta,k}^+, u_{\zeta,k}^+)_{L_\lambda^2} \end{aligned} \quad (2.12)$$

To estimate the first term in the second line of (2.12), we define $\tilde{\gamma}^+$ a cutoff function on the real line such that

$$\tilde{\gamma}^+(t) = \begin{cases} 1 & \text{if } t \in [e, e^4], \\ 0 & \text{if } t \in (-\infty, 1] \cup [e^5, +\infty). \end{cases}$$

We define $\tilde{\gamma}_k^+(\xi) = \tilde{\gamma}^+(e^{-k}\xi_{2n+1})$ then $\tilde{\gamma}_k^+ = 1$ on $\text{supp}(\gamma_k\psi^+)$ and denote by $\tilde{\Gamma}_k^+$ the pseudodifferential operator of symbol $\tilde{\gamma}_k^+$. Then we have

$$\begin{aligned} (T - e^k)e^{-\frac{\lambda}{2}}u_{\zeta,k}^+ &= (T - e^k)e^{-\frac{\lambda}{2}}\zeta\Gamma_k\Psi^+\zeta u \\ &= (T - e^k)e^{-\frac{\lambda}{2}}\zeta\tilde{\Gamma}_k^+\Gamma_k\Psi^+\zeta u \\ &= (T - e^k)\tilde{\Gamma}_k^+e^{-\frac{\lambda}{2}}\zeta\Gamma_k\Psi^+\zeta u + (T - e^k)[e^{-\frac{\lambda}{2}}\zeta, \tilde{\Gamma}_k^+]\Gamma_k\Psi^+\zeta u \\ &= (T - e^k)\tilde{\Gamma}_k^+e^{-\frac{\lambda}{2}}u_{\zeta,k}^+ + w, \end{aligned} \tag{2.13}$$

where $w := (T - e^k)[e^{-\frac{\lambda}{2}}\zeta, \tilde{\Gamma}_k^+]\Gamma_k\Psi^+\zeta u$; and hence

$$\begin{aligned} &\left(d\theta \lrcorner (T - e^k)e^{-\frac{\lambda}{2}}u_{\zeta,k}^+, e^{-\frac{\lambda}{2}}u_{\zeta,k}^+\right)_{L^2} \\ &= \left(d\theta \lrcorner (T - e^k)\tilde{\Gamma}_k^+e^{-\frac{\lambda}{2}}u_{\zeta,k}^+, e^{-\frac{\lambda}{2}}u_{\zeta,k}^+\right)_{L^2} + \left(d\theta \lrcorner w, e^{-\frac{\lambda}{2}}u_{\zeta,k}^+\right)_{L^2}. \end{aligned} \tag{2.14}$$

Since the symbol $\mathcal{S}\left((T - e^k)\tilde{\Gamma}_k^+\right) \geq 0$, we apply Lemma 2.2 for the first Levi form in the second line of (2.14) and get

$$\text{Re} \left(d\theta \lrcorner (T - e^k)\tilde{\Gamma}_k^+e^{-\frac{\lambda}{2}}u_{\zeta,k}^+, e^{-\frac{\lambda}{2}}u_{\zeta,k}^+\right)_{L^2} \geq -c\|e^{-\frac{\lambda}{2}}u_{\zeta,k}^+\|_{L^2}^2 = -c\|u_{\zeta,k}^+\|_{L_\lambda^2}^2. \tag{2.15}$$

Since $d\theta$ is a nonnegative $(1, 1)$ -form, we can use the Cauchy-Schwarz inequality for the second Levi form in the second line of (2.14) and get

$$\begin{aligned} \left|\text{Re} \left(d\theta \lrcorner w, e^{-\frac{\lambda}{2}}u_{\zeta,k}^+\right)_{L^2}\right| &\leq \sqrt{(d\theta \lrcorner w, w)(d\theta \lrcorner e^{-\frac{\lambda}{2}}u_{\zeta,k}^+, e^{-\frac{\lambda}{2}}u_{\zeta,k}^+)} \\ &\leq \frac{1}{2}e^{-k}(d\theta \lrcorner w, w)_{L^2} + \frac{1}{2}e^k \left(d\theta \lrcorner e^{-\frac{\lambda}{2}}u_{\zeta,k}^+, e^{-\frac{\lambda}{2}}u_{\zeta,k}^+\right)_{L^2} \\ &\leq ce^{-k}\|w\|_{L^2}^2 + \frac{1}{2}e^k(d\theta \lrcorner u_{\zeta,k}^+, u_{\zeta,k}^+)_{L_\lambda^2} \end{aligned} \tag{2.16}$$

where c is independent of k and λ .

From (2.12), (2.14), (2.15) and (2.16), we obtain

$$\text{Re} \left\{ \left(d\theta \lrcorner Tu_{\zeta,k}^+, u_{\zeta,k}^+\right)_{L_\lambda^2} \right\} \geq \frac{1}{2}e^k (d\theta \lrcorner u_{\zeta,k}^+, u_{\zeta,k}^+)_{L_\lambda^2} - c\|u_{\zeta,k}^+\|_{L_\lambda^2}^2 - ce^{-k}\|w\|_{L^2}^2.$$

The proof is complete by using Lemma 5.5 in Appendix to estimate $e^{-k}\|w\|^2$. Indeed,

$$\begin{aligned} e^{-k}\|w\|_{L^2}^2 &= \|(T - e^k)[e^{-\frac{\lambda}{2}}\zeta, \tilde{\Gamma}_k^+]\Gamma_k\Psi^+\zeta u\|_{L^2}^2 \\ &\leq 2e^{-k}\|T[e^{-\frac{\lambda}{2}}\zeta, \tilde{\Gamma}_k^+]\Gamma_k\Psi^+\zeta u\|_{L^2}^2 + 2e^k\|[e^{-\frac{\lambda}{2}}\zeta, \tilde{\Gamma}_k^+]\Gamma_k\Psi^+\zeta u\|_{L^2}^2 \\ &\stackrel{\text{Lemma 5.5}}{\leq} c_{s,r}e^{-k}\|\zeta u\|_{H^{-r}}^2. \end{aligned}$$

□

2.3. A special norm between $\|\zeta_0\|_{H^s}$ and $\|\zeta_1\|_{H^s}$. Let σ, ζ be cutoff functions in U such that $\sigma \prec \zeta$. Recall that $u_{\zeta,k}^\pm = \zeta\Gamma_k\Psi^\pm\zeta u$. Fix $s_o \geq 0$. Let $s \geq 0$ and $g : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that $g(t)t^{s_o+2}$ is decreasing and $g(t)t^r$ is increasing for some $r \geq s_o + 2$. We write the next several results for a generic g but we will use $g(t) = t^{-r}$ or $g(t) = t^{-r}f(t)$ in §3 to establish the a-priori estimates. We define the following spaces and norms on $(0, q)$ -forms

$$g\mathcal{A}_{\zeta;0,q}^{\pm,s\sigma} = \left\{ u \in H_{0,q}^{-s_o}(M) : \|u\|_{g\mathcal{A}_{\zeta}^{\pm,s\sigma}}^2 := \sum_{k=1}^{\infty} g^2(e^k) \|e^{ks\sigma} u_{\zeta,k}^\pm\|_{L^2}^2 < \infty \right\}.$$

Be Remark 2.3, each atom $g^2(e^k) \|e^{ks\sigma} u_{\zeta,k}^\pm\|_{L^2}^2 \leq c_{k,s}$, however, the finite of the full norm $\|u\|_{g\mathcal{A}_{\zeta}^{\pm,s\sigma}}^2$ determines the regularity of u in the H^s -spaces. In particular, by $e^{ks\sigma}\zeta \leq e^{ks}$ and $\sum_{k=0}^{\infty} \gamma_k \equiv 1$, it follows

$$\|u\|_{g\mathcal{A}_{\zeta}^{\pm,s\sigma}}^2 \leq \sum_{k=0}^{\infty} g^2(e^k) e^{2ks} \|\Gamma_k\Psi^\pm\zeta u\|_{L^2}^2 \leq c_{s,g} \|\Lambda^s g(\Lambda)\Psi^\pm\zeta u\|_{L^2}^2 \leq c_{s,g} \|g(\Lambda)\zeta u\|_{H^s}^2,$$

where the second inequality follows by Plancherel's theorem.

On the other hand, if $\zeta_0 \prec \sigma$ then

$$\|g(\Lambda)\zeta_0 u\|_{H^s}^2 \leq c_{s,g} \left(\|u\|_{g\mathcal{A}_{\zeta}^{+,s\sigma}}^2 + \|u\|_{g\mathcal{A}_{\zeta}^{-,s\sigma}}^2 + \|g(\Lambda)\Psi^0\zeta u\|_{H^s}^2 \right).$$

Thus, the norm defined by

$$\|\cdot\|_{g\mathcal{A}_{\zeta}^{s\sigma}}^2 := \|\cdot\|_{g\mathcal{A}_{\zeta}^{+,s\sigma}}^2 + \|\cdot\|_{g\mathcal{A}_{\zeta}^{-,s\sigma}}^2 + \|g(\Lambda)\Psi^0\zeta\cdot\|_{H^s}^2$$

is an intermediate norm between $\|g(\Lambda)\zeta_0\cdot\|_{H^s}^2$ and $\|g(\Lambda)\zeta\cdot\|_{H^s}^2$ for any $\zeta_0 \prec \sigma \prec \zeta$. Instead of working on the norms of H^s we will work on $\|\cdot\|_{g\mathcal{A}_{\zeta}^{\pm,s\sigma}}^2$. For convenience, we also use the following notation:

$$\begin{aligned} \|u\|_{g\partial_b\sigma\mathcal{A}_{\zeta}^{+,s\sigma}}^2 &:= \sum_{k=1}^{\infty} g^2(e^k) \|e^{ks\sigma} \partial_b\sigma \lrcorner u_{\zeta,k}^+\|_{L^2}^2, \\ \|u\|_{g\bar{\partial}_b\sigma\mathcal{A}_{\zeta}^{-,s\sigma}}^2 &:= \sum_{k=1}^{\infty} g^2(e^k) \|e^{k(s\sigma)} \bar{\partial}_b\sigma \wedge u_{\zeta,k}^-\|_{L^2}^2, \end{aligned}$$

and

$$(u, v)_{g-\mathcal{A}_\zeta^{\pm, s\sigma}} := \sum_{k=1}^{\infty} g^2(e^k)(e^{2ks\sigma} u_{\zeta, k}^{\pm}, v_{\zeta, k}^{\pm})_{L^2},$$

for $u, v \in g-\mathcal{A}_{\zeta;0,q}^{\pm, s\sigma}$. We also define the error space whose norm appears when we estimate commutators. Set

$$g-\mathcal{E}_{\zeta, \tilde{\zeta}; 0, q}^{s, b} := \{u \in H^{-s_o}(M) : \|u\|_{g-\mathcal{E}_{\zeta, \tilde{\zeta}}^{s, b}}^2 := \|g(\Lambda)\tilde{\zeta}u\|_{H^b}^2 + \|g(\Lambda)\tilde{\Psi}^0\zeta u\|_{H^{s+b}}^2 < \infty\},$$

where $\tilde{\zeta} \succ \zeta$ and the support of $\mathcal{S}(\tilde{\Psi}^0)$ belongs to $\mathcal{C}^0$. If $g(t) = t^a \tilde{g}(t)$, we write $g-\mathcal{A}_{\zeta;0,q}^{\pm, s\sigma} = \tilde{g}-\mathcal{A}_{\zeta;0,q}^{\pm, s\sigma+a}$ and $g-\mathcal{E}_{\zeta, \tilde{\zeta}; 0, q}^{s, b} = \tilde{g}-\mathcal{E}_{\zeta, \tilde{\zeta}; 0, q}^{s, a+b}$. The error norm appears when we estimate the commutators in the following lemma.

Lemma 2.5. *Let X be a differential operator of the first order. For $u \in (g-\mathcal{A}_{\zeta;0,q}^{\pm, s\sigma}) \cap (g-\mathcal{E}_{\zeta, \tilde{\zeta}; 0, q}^{s, 0})$, we have*

$$\sum_{k=1}^n g^2(e^k) \|e^{ks\sigma} [\zeta \Gamma_k \Psi^{\pm} \zeta, X] u\|_{L^2}^2 \leq c_{s, g} \left(\|u\|_{g-\mathcal{A}_\zeta^{\pm, s\sigma}}^2 + \|u\|_{g-\mathcal{E}_{\zeta, \tilde{\zeta}}^{s, 0}}^2 \right).$$

Proof. Multiplying with $g^2(e^k)$ to both sides of the estimate in Lemma 5.6(i) in Appendix, and then taking the summation over k , we get the desired estimate. $\square$

The combination of Lemma 2.5 and 5.6(ii) in Appendix gives us

$$\|Xu\|_{g-\mathcal{A}_\zeta^{\pm, s\sigma}}^2 \leq c_{s, g} \left(\|u\|_{g-\mathcal{A}_\zeta^{\pm, s\sigma+1}}^2 + \|u\|_{g-\mathcal{E}_{\zeta, \tilde{\zeta}}^{s, 1}}^2 \right),$$

for $u \in (g-\mathcal{A}_{\zeta;0,q}^{\pm, s\sigma+1}) \cap (g-\mathcal{E}_{\zeta, \tilde{\zeta}; 0, q}^{s, 1})$. Consequently, we have the following corollary.

Corollary 2.6. (i) *If $u \in (g-\mathcal{A}_{\zeta;0,q}^{\pm, s\sigma+1}) \cap (g-\mathcal{E}_{\zeta, \tilde{\zeta}; 0, q}^{s, 1})$ then $\bar{\partial}_b u, \bar{\partial}_b^* u \in (g-\mathcal{A}_{\zeta;0,q}^{\pm, s\sigma}) \cap (g-\mathcal{E}_{\zeta, \tilde{\zeta}; 0, q}^{s, 0})$*

(ii) *If $u \in (g-\mathcal{A}_{\zeta;0,q}^{\pm, s\sigma+2}) \cap (g-\mathcal{E}_{\zeta, \tilde{\zeta}; 0, q}^{s, 2})$ then $\bar{\partial}_b^* \bar{\partial}_b u, \bar{\partial}_b \bar{\partial}_b^* u, \square_b u, \square_b^\delta u \in (g-\mathcal{A}_{\zeta;0,q}^{\pm, s\sigma}) \cap (g-\mathcal{E}_{\zeta, \tilde{\zeta}; 0, q}^{s, 0})$, where $\square_b^\delta$ is an elliptic perturbation of $\square_b$ defined in §3.3.*

Note on notation. In what follows we use notation $A \lesssim B$ for $A \leq cB$ where c is a constant depending on indices and cutoff functions of the space $\|\cdot\|_{g-\mathcal{A}_{\zeta;0,q}^{\pm, s\sigma+a}}$ and $\|\cdot\|_{g-\mathcal{E}_{\zeta, \tilde{\zeta}; 0, q}^{s\sigma, a}}$ such as $s, s_o, a, g, \zeta, \sigma$ but not on k .

3. A-PRIORI ESTIMATES

In this section we establish the *a-priori* estimates for the system $(\bar{\partial}_b, \bar{\partial}_b^*)$, $\square_b$ and $\square_b^\delta$ which are needed in the proofs of the main theorems.

3.1. A-priori estimates for the system $(\bar{\partial}_b, \bar{\partial}_b^*)$. The central of this subsection is to prove the following *a-priori* estimates for the system $(\bar{\partial}_b, \bar{\partial}_b^*)$ in the special norm $\|\cdot\|_{g-\mathcal{A}_\zeta^{\pm, s\sigma}}$. The proof of Theorem 1.2 is also given at the end of this subsection.

Theorem 3.1. *Suppose that the σ -superlogarithmic property on $(0, q_0)$ -forms holds with the pair of rates (f, f_0) . Then the following holds:*

(i) *For $q \geq q_0$, we have*

$$\|u\|_{gf-\mathcal{A}_\zeta^+, s\sigma}^2 + \|u\|_{gf_0 \ln -\partial_b \sigma \lrcorner \mathcal{A}_\zeta^+, s\sigma}^2 \leq c \left(\|\bar{\partial}_b u\|_{g-\mathcal{A}_\zeta^+, s\sigma}^2 + \|\bar{\partial}_b^* u\|_{g-\mathcal{A}_\zeta^+, s\sigma}^2 \right) + c_{s,g} \|u\|_{g-\mathcal{E}_{\zeta, \tilde{\zeta}}^{s,0}}^2. \quad (3.1)$$

for all $u \in (g-\mathcal{A}_{\zeta;0,q}^{+, s\sigma+1}) \cap (g-\mathcal{E}_{\zeta, \tilde{\zeta};0,q}^{s,1})$.

(ii) *For $q \leq n - q_0$, we have*

$$\|u\|_{gf-\mathcal{A}_\zeta^-, s\sigma}^2 + \|u\|_{gf_0 \ln -\bar{\partial}_b \sigma \wedge \mathcal{A}_\zeta^-, s\sigma}^2 \leq c \left(\|\bar{\partial}_b u\|_{g-\mathcal{A}_\zeta^-, s\sigma}^2 + \|\bar{\partial}_b^* u\|_{g-\mathcal{A}_\zeta^-, s\sigma}^2 \right) + c_{s,g} \|u\|_{g-\mathcal{E}_{\zeta, \tilde{\zeta}}^{s,0}}^2. \quad (3.2)$$

for all $u \in (g-\mathcal{A}_{\zeta;0,q}^-, s\sigma+1) \cap (g-\mathcal{E}_{\zeta, \tilde{\zeta};0,q}^{s,1})$.

The proof of this theorem is divided into four steps:

- Step 1. We point out that if the σ -superlogarithmic property holds on $(0, q_0)$ forms then it still holds on $(0, q)$ -forms with $q_0 \leq q \leq n$.
- Step 2. Using the Kohn-Morrey-Hörmander inequality in Proposition 2.1 and the estimate of the Levi form $(d\theta \lrcorner T \cdot, \cdot)_{L_\lambda^2}$ in Proposition 2.4, we obtain the estimate for each atom $\|e^{ks\sigma} u_{\zeta,k}^\pm\|_{L^2}$ of the full norm $\|\cdot\|_{g-\mathcal{A}_\zeta^{\pm, s\sigma}}$. Indeed,

$$\begin{aligned} f^2(e^k) \|e^{ks\sigma} u_{\zeta,k}^+\|_{L^2}^2 + k^2 f_0^2(e^k) \|e^{ks\sigma} \partial_b \sigma \lrcorner u_{\zeta,k}^+\|_{L^2}^2 \\ \leq c \left(\|e^{ks\sigma} \bar{\partial}_b u_{\zeta,k}^+\|_{L^2}^2 + \|e^{ks\sigma} \bar{\partial}_b^* u_{\zeta,k}^+\|_{L^2}^2 \right) + c_{s,r} e^{-k} \|\zeta u\|_{H^{-r}}^2 \end{aligned} \quad (3.3)$$

for all $u \in H_{0,q}^{-s_0}(M)$, $s, r \in \mathbb{R}^+$ with $r \geq s_0$ and $q \geq q_0$.

- Step 3. A similar estimate for negative microlocalization is obtained by the aid of Hodge-* on complementary degree. In particular, (3.3) is equivalent to

$$\begin{aligned} f^2(e^k) \|e^{ks\sigma} u_{\zeta,k}^-\|_{L^2}^2 + k^2 f_0^2(e^k) \|e^{ks\sigma} \bar{\partial}_b \wedge u_{\zeta,k}^-\|_{L^2}^2 \\ \leq c \left(\|e^{ks\sigma} \bar{\partial}_b u_{\zeta,k}^-\|_{L^2}^2 + \|e^{ks\sigma} \bar{\partial}_b^* u_{\zeta,k}^-\|_{L^2}^2 \right) + c_{s,r} e^{-k} \|\zeta u\|_{H^{-r}}^2 \end{aligned} \quad (3.4)$$

for all $u \in H_{0,q}^{-s_0}(M)$, $s, r \in \mathbb{R}^+$ with $r \geq s_0$ and $q \leq n - q_0$.

- Step 4. Multiplying with $g^2(e^k)$ to both sides of (3.3) and (3.4), and then taking summation over k , we get the desired estimates (3.1) and (3.2), respectively.

The proof of Step 1 immediately follows by following proposition.

Proposition 3.2. *If the σ -superlogarithmic property holds on $(0, q)$ -forms then it still holds on $(0, q+1)$ -forms with the same rates.*

Remark 3.3. Let $\mathcal{M}$ be a nonnegative $(1, 1)$ -form. It is obvious that the $(f\text{-}\mathcal{M})$ -property on $(0, q)$ -forms implies the $(f\text{-}\mathcal{M})$ -property on $(0, q+1)$ -forms. However, in the definition of the σ -superlogarithmic property, the Levi form $\mathcal{L}_\sigma$ is not a nonnegative $(1, 1)$ -form in general. So we have to give a direct proof for Proposition 3.2.

Proof. We assume that there exist functions $f, f_0 : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with $f, f_0 \gg 1$ and a family of smooth real-valued functions $\{\lambda_t\}_{t \geq 1}$ defined on U such that

$$\begin{cases} \langle (\mathcal{L}_{\lambda_t} + td\theta) \lrcorner v, v \rangle \geq f_0^2(t) (\ln t |\langle \mathcal{L}_\sigma \lrcorner v, v \rangle| + \ln^2 t |\partial_b \sigma \lrcorner v|^2) + f^2(t) |v|^2, \\ |T^m(\lambda_t)| \leq c_m t^{\frac{m}{2}}, \text{ for all } m \in \mathbb{N}, \end{cases} \quad (3.5)$$

holds at any $x \in U$ for all smooth $(0, q)$ -forms v . Let $u = \sum_{L \in \mathcal{I}_{q+1}} u_L \bar{\omega}_L$ is a $(0, q+1)$ -form defined in U . We rewrite u as a non-ordered sum

$$u = \frac{1}{(q+1)!} \sum_{|L|=q+1} u_L \bar{\omega}_L = \frac{(-1)^q}{q+1} \sum_{l=1}^n \left(\frac{1}{q!} \sum_{|J|=q} u_{lJ} \bar{\omega}_J \right) \wedge \bar{\omega}_l.$$

For $l = 1, \dots, n$, we define a set of $(0, q)$ -forms v_l by

$$v_l := \frac{1}{q!} \sum_{|J|=q} u_{lJ} \bar{\omega}_J = \sum_{J \in \mathcal{I}_q} u_{lJ} \bar{\omega}_J.$$

It is easy to check that

$$\begin{cases} \sum_{l=1}^n |v_l|^2 = (q+1) |u|^2; \\ \sum_{l=1}^n |\lambda \lrcorner v_l|^2 = q |\lambda \lrcorner u|^2 \text{ if } \lambda \text{ is a } (1, 0)\text{-form}; \\ \sum_{l=1}^n \langle \lambda \lrcorner v_l, v_l \rangle = q \langle \lambda \lrcorner u, u \rangle \text{ if } \lambda \text{ is a } (1, 1)\text{-form}. \end{cases}$$

Thus,

$$\begin{aligned} & \sum_{l=1}^n \left(f_0^2(t) (\ln t |\langle \mathcal{L}_\sigma \lrcorner v_l, v_l \rangle| + \ln^2 t |\partial_b \sigma \lrcorner v_l|^2) + f^2(t) |v_l|^2 \right) \\ &= q f_0^2(t) (\ln t |\langle \mathcal{L}_\sigma \lrcorner u, u \rangle| + \ln^2 t |\partial_b \sigma \lrcorner u|^2) + (q+1) f^2(t) |u|^2. \end{aligned} \quad (3.6)$$

Substituting $v = v_l$ into (3.5) and taking summation over l , to get

$$\begin{aligned} & \sum_{l=1}^n \left(f_0^2(t) (\ln t |\langle \mathcal{L}_\sigma \lrcorner v_l, v_l \rangle| + \ln^2 t |\partial_b \sigma \lrcorner v_l|^2) + f^2(t) |v_l|^2 \right) \\ & \leq \sum_{l=1}^n \left(\langle (\mathcal{L}_{\lambda_t} + td\theta) \lrcorner v_l, v_l \rangle \right) = q \langle (\mathcal{L}_{\lambda_t} + td\theta) \lrcorner u, u \rangle. \end{aligned} \quad (3.7)$$

From (3.6) and (3.7), the σ -superlogarithmic property on $(0, q+1)$ -forms holds. $\square$

The second step in the proof of Theorem 3.1 is contained in the following proposition.

Proposition 3.4. *Suppose that the σ -superlogarithmic property with the pair of rates (f, f_0) holds on $(0, q)$ -forms. Then*

$$\begin{aligned} f^2(e^k) \|e^{ks\sigma} u_{\zeta,k}^+\|_{L^2}^2 + k^2 f_0^2(e^k) \|\partial_b \sigma \lrcorner e^{ks\sigma} u_{\zeta,k}^+\|_{L^2}^2 \\ \leq c \left(\|e^{ks\sigma} \bar{\partial}_b u_{\zeta,k}^+\|_{L^2}^2 + \|e^{ks\sigma} \bar{\partial}_b^* u_{\zeta,k}^+\|_{L^2}^2 \right) + c_{s,r} e^{-k} \|\zeta u\|_{H^{-r}}^2, \end{aligned} \quad (3.8)$$

for all $u \in H_{0,q}^{-s_0}(M)$, $k = 1, 2, \dots$, and $s, r \in \mathbb{R}$ with $r \geq s_0$, where c and $c_{s,r}$ are independent of k .

Proof. Assume that for any $t \geq 1$ there exists a real-valued function λ_t smooth in a neighborhood of $\text{supp}(\sigma)$ such that

$$\begin{cases} \langle (\mathcal{L}_{\lambda_t} + td\theta) \lrcorner u, u \rangle \geq f_0^2(t) (\ln t |\langle \mathcal{L}_{\sigma} \lrcorner u, u \rangle| + \ln^2 t |\partial_b \sigma \lrcorner u|^2) + f^2(t) |u|^2, \\ |T^m(\lambda_t)| \leq c_m t^{\frac{m}{2}} \text{ for all } m \in \mathbb{N}, \end{cases} \quad (3.9)$$

for all $u \in C_{0,q}^\infty(M)$. We are going to apply Proposition 2.1 and 2.4 for

$$\lambda := \lambda_{k,s} := \chi(\lambda_{e^k}) - 2ks\sigma$$

where $\chi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a smooth function and will be chosen later. By the definition of $\mathcal{L}_\lambda$, we have

$$\mathcal{L}_\lambda = \dot{\chi} \mathcal{L}_{\lambda_{e^k}} + \ddot{\chi} \partial_b(\lambda_{e^k}) \wedge \bar{\partial}_b(\lambda_{e^k}) - 2sk \mathcal{L}_\sigma,$$

and hence

$$\langle \mathcal{L}_\lambda \lrcorner u, u \rangle = \dot{\chi} \langle \mathcal{L}_{\lambda_{e^k}} \lrcorner u, u \rangle + \ddot{\chi} |\partial_b \lambda_{e^k} \lrcorner u|^2 - 2sk \langle \mathcal{L}_\sigma \lrcorner u, u \rangle. \quad (3.10)$$

On other hand, it follows from (2.9),

$$\bar{\partial}_{b,\lambda}^* u = \bar{\partial}_b^* u - \partial_b \lambda \lrcorner u = \bar{\partial}_b^* u - \dot{\chi} \partial_b \lambda_{e^k} \lrcorner u + 2sk \partial_b \sigma \lrcorner u. \quad (3.11)$$

Thus we get from Proposition 2.1 applied to $u_{\zeta,k}^+ \in C_{0,q}^\infty(M)$ supported in U (by Remark 2.3), under the choice of the weight $\lambda := \lambda_{k,s}$ and taking into account (3.10) and (3.11),

$$\begin{aligned} \|e^{-\frac{\lambda}{2}} \bar{\partial}_b u_{\zeta,k}^+\|_{L^2}^2 + 3 \|e^{-\frac{\lambda}{2}} \bar{\partial}_b^* u_{\zeta,k}^+\|_{L^2}^2 + c \|e^{-\frac{\lambda}{2}} u_{\zeta,k}^+\|_{L^2}^2 \\ \geq \text{Re} \int_M [\dot{\chi} \langle \mathcal{L}_{\lambda_{e^k}} \lrcorner u_{\zeta,k}^+, u_{\zeta,k}^+ \rangle + \ddot{\chi} |\partial_b \lambda_{e^k} \lrcorner u_{\zeta,k}^+|^2 + \langle d\theta \lrcorner T u_{\zeta,k}^+, u_{\zeta,k}^+ \rangle] e^{-\lambda} dV \\ - \int_M [2sk \langle \mathcal{L}_\sigma \lrcorner u_{\zeta,k}^+, u_{\zeta,k}^+ \rangle + 3\dot{\chi}^2 |\partial_b \lambda_{e^k} \lrcorner u_{\zeta,k}^+|^2 + 12s^2 k^2 |\partial_b \sigma \lrcorner u_{\zeta,k}^+|^2] e^{-\lambda} dS. \end{aligned} \quad (3.12)$$

Since λ_{e^k} is uniformly bounded, we can assume that $-1 \leq \lambda_{e^k} \leq 0$. We choose $\chi(a) = \frac{1}{3}e^a$. It follows: (i) $\ddot{\chi} \geq 3\dot{\chi}^2$ so we can remove the term involving $|\partial_b \lambda_{e^k} \lrcorner u_{\zeta,k}^+|^2$; (ii) by the second line of 3.9 it follows

$$|\lambda| \leq c_{0,s} k \quad \text{and} \quad |T^m(\lambda)| \leq c_{m,s} e^{\frac{mk}{2}},$$

so we can use Proposition 2.4 for to estimate the Levi form $\langle d\theta \lrcorner T \cdot, \cdot \rangle$ as

$$\operatorname{Re} \int_M \langle d\theta \lrcorner T u_{\zeta,k}^+, u_{\zeta,k}^+ \rangle e^{-\lambda} dV \geq \frac{1}{2} e^k \int_M \langle d\theta \lrcorner u_{\zeta,k}^+, u_{\zeta,k}^+ \rangle e^{-\lambda} dV - c \|e^{\frac{\lambda}{2}} u_{\zeta,k}^+\|^2 - c_{r,s} e^{-k} \|\zeta u\|_{H^{-r}}^2;$$

(iii) $e^{-\chi} \approx 1$ and $\dot{\chi} \gtrsim 1$, so we remove the factor $e^{-\chi}$ in $e^{-\lambda} = e^{-\chi+2ks\sigma}$. We can conclude that

$$\begin{aligned} & \|e^{sk\sigma} \bar{\partial}_b u_{\zeta,k}^+\|_{L^2}^2 + \|e^{sk\sigma} \bar{\partial}_b^* u_{\zeta,k}^+\|_{L^2}^2 + c_{s,r} e^{-k} \|\zeta u\|_{H^{-r}}^2 \\ & \geq c \int_M \left[\langle \mathcal{L}_{\lambda_{e^k} \lrcorner} u_{\zeta,k}^+, u_{\zeta,k}^+ \rangle + \frac{1}{2} e^k \langle d\theta \lrcorner u_{\zeta,k}^+, u_{\zeta,k}^+ \rangle \right] e^{2sk\sigma} dV \\ & \quad - C \int_M [|u_{\zeta,k}^+|^2 + 12s^2 k^2 |\partial_b \sigma \lrcorner u_{\zeta,k}^+|^2 + sk \langle \mathcal{L}_{\sigma \lrcorner} u_{\zeta,k}^+, u_{\zeta,k}^+ \rangle] e^{2sk\sigma} dV \\ & \geq c' \left(f^2(e^k) \|e^{ks\sigma} u_{\zeta,k}^+\|_{L^2}^2 + k^2 f_0^2(e^k) \|\partial_b \sigma \lrcorner e^{ks\sigma} u_{\zeta,k}^+\|_{L^2}^2 + k f_0^2(e^k) \int_M |\langle \mathcal{L}_{\sigma \lrcorner} u_{\zeta,k}^+, u_{\zeta,k}^+ \rangle| e^{2ks\sigma} dV \right) \\ & \quad - C \int_M [|u_{\zeta,k}^+|^2 + 12s^2 k^2 |\partial_b \sigma \lrcorner u_{\zeta,k}^+|^2 + sk \langle \mathcal{L}_{\sigma \lrcorner} u_{\zeta,k}^+, u_{\zeta,k}^+ \rangle] e^{2sk\sigma} dV. \end{aligned} \tag{3.13}$$

where the last inequality follows the first line of (3.9) for $t = e^k$. We obtain the desired inequality since the last line is absorbed by the line before for large k since both $f, f_0 \gg 1$, otherwise it is absorbed by $c_{s,r} e^{-k} \|\zeta u\|_{H^{-r}}^2$. $\square$

The estimates for positive and negative microlocalizations are related, in complementary degree, by the aid of the Hodge- $*$ theory.

Proposition 3.5. *Fix $1 \leq q \leq n$ and $r \geq s_o$. The estimates*

$$\begin{aligned} & c_{s,r} e^{-k} \|\zeta u\|_{H^{-r}}^2 + c \left(\|e^{ks\sigma} \bar{\partial}_b u_{\zeta,k}^+\|_{L^2}^2 + \|e^{ks\sigma} \bar{\partial}_b^* u_{\zeta,k}^+\|_{L^2}^2 \right) \\ & \geq f^2(e^k) \|e^{ks\sigma} u_{\zeta,k}^+\|_{L^2}^2 + f_0^2(e^k) k^2 \|\partial_b \sigma \lrcorner e^{ks\sigma} u_{\zeta,k}^+\|_{L^2}^2 \end{aligned} \tag{3.14}$$

for all $u \in H_{0,q}^{-s_o}(M)$ and

$$\begin{aligned} & c_{s,r} e^{-k} \|\zeta u\|_{H^{-r}}^2 + c \left(\|e^{ks\sigma} \bar{\partial}_b u_{\zeta,k}^-\|_{L^2}^2 + \|e^{ks\sigma} \bar{\partial}_b^* u_{\zeta,k}^-\|_{L^2}^2 \right) \\ & \geq f^2(e^k) \|e^{ks\sigma} u_{\zeta,k}^-\|_{L^2}^2 + f_0^2(e^k) k^2 \|\bar{\partial}_b \sigma \wedge e^{ks\sigma} u_{\zeta,k}^-\|_{L^2}^2 \end{aligned} \tag{3.15}$$

for all $u \in H_{0,n-q}^{-s_o}(M)$ are equivalent.

Proof. We only need to prove (3.14) $\implies$ (3.15) since the reversed direction follows analogously. We define the local conjugate-linear duality map $F^{n-q} : C_{0,n-q}^\infty(M) \rightarrow C_{0,q}^\infty(M)$ as follows. If $u = \sum'_{|J|=n-q} u_J \bar{\omega}_J$ then

$$F^{n-q} u = \sum \epsilon_{\{1,\dots,n\}}^{\{J,J'\}} \bar{u}_J \bar{\omega}_{J'},$$

where J' denotes the strictly increasing q -tuple consisting of all integers in $[1, n]$ which do not belong to J and $\epsilon_{\{1, n\}}^{J, J'}$ is the sign of the permutation $\{J, J'\} \sim \{1, \dots, n\}$.

By this definition, $F^q F^{n-q} u = u$, $\|e^{ks\sigma} F^{n-q} u\| = \|e^{ks\sigma} u\|$ and furthermore

$$\begin{cases} \bar{\partial}_b F^{n-q} u = F^{n-q-1} (\bar{\partial}_b^* u + \dots), \\ \bar{\partial}_b^* F^{n-q} u = F^{n-q+1} (\bar{\partial}_b u + \dots), \\ \partial_b \sigma \lrcorner F^{n-q} u = F^{n-q+1} (\bar{\partial}_b \sigma \wedge u), \end{cases}$$

for any $(0, n-q)$ -form u supported in U , where dots refers the term in which u is not differentiated. It follows

$$\begin{cases} \|e^{ks\sigma} \bar{\partial}_b F^{n-q} u\|^2 + \|e^{ks\sigma} \bar{\partial}_b^* F^{n-q} u\|^2 \leq 2(\|e^{ks\sigma} \bar{\partial}_b u\|^2 + \|e^{ks\sigma} \bar{\partial}_b^* u\|^2) + C\|e^{ks\sigma} u\|^2 \\ \|e^{ks\sigma} F^{n-q} u\| + k^2 \|e^{ks\sigma} \partial_b \sigma \lrcorner F^{n-q} u\| = \|e^{ks\sigma} u\| + \|e^{ks\sigma} \bar{\partial}_b \sigma \wedge u\|^2. \end{cases} \quad (3.16)$$

On the other hand for each coefficient $u_{k,J}^-$ of $u_{\zeta,k}^-$ we have

$$\begin{aligned} \overline{u_{k,J}^-}(x) &= \overline{(\zeta \Gamma_k^- \Psi_k^- \zeta u_J)(x)} \\ &= (2\pi)^{2n+1} \zeta(x) \overline{\int_{\mathbb{R}_\xi^{2n+1}} e^{ix\xi} \gamma_k(\xi) \psi^-(\xi) \int_{\mathbb{R}_y^{2n+1}} e^{-iy\xi} (\zeta u_J)(y) dy d\xi} \\ &= (2\pi)^{2n+1} \zeta(x) \int_{\mathbb{R}_\xi^{2n+1}} e^{-ix\xi} \gamma(-e^k \xi_{2n+1}) \psi(-\xi) \int_{\mathbb{R}_y^{2n+1}} e^{iy\xi} \overline{(\zeta u_J)(y)} dy d\xi \\ &\stackrel{\xi := -\xi}{=} -(-2\pi)^{2n+1} \zeta(x) \int_{\mathbb{R}_\xi^{2n+1}} e^{ix\xi} \gamma(e^k \xi_{2n+1}) \psi(\xi) \int_{\mathbb{R}_y^{2n+1}} e^{-iy\xi} \zeta(y) \overline{u_J(y)} dy d\xi \\ &= -(-2\pi)^{2n+1} \zeta(x) \int_{\mathbb{R}_\xi^{2n+1}} e^{ix\xi} \gamma_k(\xi_{2n+1}) \psi^+(\xi) \int_{\mathbb{R}_y^{2n+1}} e^{-iy\xi} (\zeta \bar{u}_J)(y) dy d\xi \\ &= -(\zeta \Gamma_k^+ \Psi^+ \zeta \bar{v})(x) = -\bar{u}_{k,J}^+(x), \end{aligned} \quad (3.17)$$

and hence $F^{n-q}(u_{\zeta,k}^-) = -(\overline{F^{n-q} u})_k^+$ is a positive microlocalized $(0, q)$ -form supported on U . So we apply (3.14) for $F^{n-q}(u_{\zeta,k}^-)$ together with (3.16) for u replaced by $u_{\zeta,k}^-$, we get (3.15). The proof of Proposition 3.5 is complete. $\square$

Now we are ready to get the complete the proof of Theorem 3.1.

End proof of Theorem 3.1. Using the estimate (3.3) for $k = 1, 2, \dots$ with the choice of r such that $t^{-r} \leq g(t)$ for all t , we multiply by $g^2(e^k)$ and sum over k to establish

$$\begin{aligned}
& \|u\|_{gf-\mathcal{A}_\zeta^+, s\sigma}^2 + \|u\|_{gf_0 \ln -\partial_b \sigma \lrcorner \mathcal{A}_\zeta^+, s\sigma}^2 \\
&= \sum_{k=1}^n g^2(e^k) \left(f^2(e^k) \|e^{ks\sigma} u_{\zeta,k}^+\|_{L^2}^2 + f_0^2(e^k) k^2 \|\partial \sigma \lrcorner e^{ks\sigma} u_{\zeta,k}^+\|_{L^2}^2 \right) \\
&\leq c \sum_{k=1}^n g^2(e^k) \left(\|e^{ks\sigma} \bar{\partial}_b u_{\zeta,k}^+\|_{L^2}^2 + \|e^{ks\sigma} \bar{\partial}_b^* u_{\zeta,k}^+\|_{L^2}^2 + c_{s,g} e^{-k} \|g(\Lambda) \zeta u\|_{L^2}^2 \right) \\
&\leq c \sum_{k=1}^n g^2(e^k) \left(\|e^{ks\sigma} (\bar{\partial}_b u)_{\zeta,k}^+\|_{L^2}^2 + \|e^{ks\sigma} (\bar{\partial}_b^* u)_{\zeta,k}^+\|_{L^2}^2 \right. \\
&\quad \left. + \|e^{ks\sigma} [\zeta \Gamma_k \Psi^+ \zeta, \bar{\partial}_b] u\|_{L^2}^2 + \|e^{ks\sigma} [\zeta \Gamma_k \Psi^+ \zeta, \bar{\partial}_b^*] u\|_{L^2}^2 + c_{s,g} e^{-k} \|g(\Lambda) \zeta u\|_{L^2}^2 \right) \\
&\leq c \left(\|\bar{\partial}_b u\|_{g-\mathcal{A}_\zeta^+, s\sigma}^2 + \|\bar{\partial}_b^* u\|_{g-\mathcal{A}_\zeta^+, s\sigma}^2 \right) + c_{s,g} \left(\|u\|_{g-\mathcal{A}_\zeta^+, s\sigma}^2 + \|u\|_{g-\mathcal{E}_{\zeta,\bar{\zeta}}^{s,0}}^2 \right),
\end{aligned} \tag{3.18}$$

where the last inequality follows by Lemma 2.5. Since $f \gg 1$, the term $c_{s,g} \|u\|_{g-\mathcal{A}_\zeta^+, s\sigma}^2$ can be absorbed by $\|u\|_{gf-\mathcal{A}_\zeta^+, s\sigma}^2$ in the first line of (3.18). Then the desired estimate (3.3) follows. The negative microlocal estimate (3.4) is proved analogously. $\square$

We also have a proof of Theorem 1.2 as a consequence of Remark 3.3 and Propositions 3.4 and 3.5 as follows.

Proof of Theorem 1.2. By Remark 3.3 the $(f-\mathcal{M})$ property holds on $(0, q)$ -forms with $q \geq q_0$. Similar to the argument in Proposition 3.4 but with simpler calculation by the choice of weight $\lambda = \chi(\lambda_{e^k})$, we get the $(f-\mathcal{M})$ -estimate for each $u_{\zeta,k}^+$ as

$$f(e^k)^2 (\mathcal{M} \lrcorner u_{\zeta,k}^+, u_{\zeta,k}^+)_{L^2} \lesssim \|\bar{\partial}_b u_{\zeta,k}^+\|_{L^2}^2 + \|\bar{\partial}_b^* u_{\zeta,k}^+\|_{L^2}^2 + \|u_{\zeta,k}^+\|_{L^2}^2 + e^{-k} \|\zeta u\|_{H^{-1}}^2. \tag{3.19}$$

for $u \in C_{0,q}^\infty(M)$. In order to get the estimate for full $u_\zeta^+ := \zeta \Psi^+ \zeta u$, we first decompose $u_\zeta^+ = \sum_{k=0}^\infty u_{\zeta,k}^+$ and hence

$$(\mathcal{M} \lrcorner f(\Lambda) u_\zeta^+, f(\Lambda) u_\zeta^+)_{L^2} = \sum_{k, \hat{k}} (\mathcal{M} \lrcorner f(\Lambda) u_{\zeta,k}^+, f(\Lambda) u_{\zeta,\hat{k}}^+)_{L^2}. \tag{3.20}$$

We first consider the case $|k - \hat{k}| \geq 3$. In this case we have $\text{supp}(\gamma_k) \cap \text{supp}(\gamma_{\hat{k}}) = \emptyset$ and hence, by the argument in Lemma 5.5 below we can get

$$\left| (\mathcal{M} \lrcorner f(\Lambda) u_{\zeta,k}^+, f(\Lambda) u_{\zeta,\hat{k}}^+)_{L^2} \right| \lesssim e^{-(k+\hat{k})} \|\zeta u\|_{H^{-1}}^2 \quad \text{for all } |k - \hat{k}| \geq 3.$$

Otherwise for $|k - \hat{k}| \leq 3$, we use the Cauchy-Schwarz inequality since the Hermitian form $(\mathcal{M}_\perp, \cdot)$ is nonnegative. Therefore we obtain

$$\begin{aligned}
(\mathcal{M}_\perp f(\Lambda)u_\zeta^+, f(\Lambda)u_\zeta^+)_{L^2} &\lesssim \sum_{k=1}^{\infty} \left((\mathcal{M}_\perp f(\Lambda)u_{\zeta,k}^+, f(\Lambda)u_{\zeta,k}^+)_{L^2} + e^{-k} \|\zeta u\|_{H^{-1}}^2 \right) \\
&\stackrel{\text{Gårding's inequality}}{\lesssim} \sum_{k=1}^{\infty} \left(f^2(e^k) (\mathcal{M}_\perp u_{\zeta,k}^+, u_{\zeta,k}^+)_{L^2} + \|u_{\zeta,k}^+\|_{L^2}^2 + e^{-k} \|\zeta u\|_{H^{-1}}^2 \right) \\
&\stackrel{\text{by (3.19)}}{\lesssim} \sum_{k=1}^{\infty} \left(\|\bar{\partial}_b u_{\zeta,k}^+\|_{L^2}^2 + \|\bar{\partial}_b^* u_{\zeta,k}^+\|_{L^2}^2 + \|u_{\zeta,k}^+\|_{L^2}^2 + e^{-k} \|\zeta u\|_{-1}^2 \right) \\
&\stackrel{\text{by Lemma 2.5}}{\lesssim} \|\bar{\partial}_b u_\zeta^+\|_{L^2}^2 + \|\bar{\partial}_b^* u_\zeta^+\|_{L^2}^2 + \|\tilde{\zeta} u\|_{L^2}^2
\end{aligned} \tag{3.21}$$

for all $u \in C_{0,q}^\infty(M)$, where $\tilde{\zeta} \succ \zeta$. By [Kha16, Theorem 5] (or analogous proof of Proposition 3.5), (3.21) is equivalent to

$$\|\sqrt{\text{Tr}(\mathcal{M})} \times f(\Lambda)u_\zeta^-\|_{L^2}^2 - (\mathcal{M}_\perp f(\Lambda)u_\zeta^-, f(\Lambda)u_\zeta^-)_{L^2} \lesssim \|\bar{\partial}_b u_\zeta^-\|_{L^2}^2 + \|\bar{\partial}_b^* u_\zeta^-\|_{L^2}^2 + \|\tilde{\zeta} u\|_{L^2}^2. \tag{3.22}$$

for all $u \in C_{0,q}^\infty(M)$ with $q \leq n - q_0$. This is complete the proof of Theorem 1.2. $\square$

3.2. A-priori estimates for $\square_b$. We need technique lemmas before going estimates for $\square_b$ in our norms.

Lemma 3.6. *Let $a \geq 0$ and $\epsilon > 0$. Then, there exists c_ϵ such that the following holds.*

(i) *for $v \in (g\mathcal{A}_{\zeta;0,q-1}^{+,s\sigma+a}) \cap (g\mathcal{E}_{\zeta,\tilde{\zeta};0,q-1}^{s,a})$ and $w \in (g\mathcal{A}_{\zeta;0,q}^{+,s\sigma+1-a}) \cap (g\mathcal{E}_{\zeta,\tilde{\zeta};0,q}^{s,-a})$, we have*

$$\begin{aligned}
\left| \text{Re} \left\{ (v, \bar{\partial}_b^* w)_{g\mathcal{A}_\zeta^{+,s\sigma}} - (\bar{\partial}_b v, w)_{g\mathcal{A}_\zeta^{+,s\sigma}} \right\} \right| &\leq \epsilon \left(\|v\|_{g\mathcal{A}_\zeta^{+,s\sigma+a}}^2 + \|v\|_{g\mathcal{E}_{\zeta,\tilde{\zeta}}^{s,a}}^2 \right) \\
&+ c_\epsilon \left(\|w\|_{g\mathcal{A}_\zeta^{+,s\sigma-a}}^2 + \|w\|_{g\ln - \bar{\partial}_b \sigma \wedge \mathcal{A}_\zeta^{+,s\sigma-a}}^2 + \|w\|_{g\mathcal{E}_{\zeta,\tilde{\zeta}}^{s,-a}}^2 \right)
\end{aligned} \tag{3.23}$$

(ii) *For $v \in (g\mathcal{A}_{\zeta;0,q-1}^{-,s\sigma+a}) \cap (g\mathcal{E}_{\zeta,\tilde{\zeta};0,q-1}^{s,a})$ and $w \in (g\mathcal{A}_{\zeta;0,q}^{-,s\sigma+1-a}) \cap (g\mathcal{E}_{\zeta,\tilde{\zeta};0,q}^{s,-a})$, we have*

$$\begin{aligned}
\left| \text{Re} \left\{ (v, \bar{\partial}_b w)_{g\mathcal{A}_\zeta^{-,s\sigma}} - (\bar{\partial}_b^* v, w)_{g\mathcal{A}_\zeta^{-,s\sigma}} \right\} \right| &\leq \epsilon \left(\|v\|_{g\mathcal{A}_\zeta^{-,s\sigma+a}}^2 + \|v\|_{g\mathcal{E}_{\zeta,\tilde{\zeta}}^{s,a}}^2 \right) \\
&+ c_\epsilon \left(\|w\|_{g\mathcal{A}_\zeta^{-,s\sigma-a}}^2 + \|w\|_{g\ln - \bar{\partial}_b \sigma \wedge \mathcal{A}_\zeta^{-,s\sigma-a}}^2 + \|w\|_{g\mathcal{E}_{\zeta,\tilde{\zeta}}^{s,-a}}^2 \right).
\end{aligned} \tag{3.24}$$

Proof. We first work on each atom of the inner product $(\cdot, \cdot)_{g-\mathcal{A}_\zeta^+, s\sigma}$,

$$\begin{aligned}
& |(e^{2ks\sigma} v_{\zeta,k}^+, (\bar{\partial}_b^* w)_{\zeta,+}^+)_{L^2} - (e^{2ks\sigma} (\bar{\partial}_b v)_{\zeta,k}^+, w_{\zeta,k}^+)_{L^2}| \\
&= |(e^{2ks\sigma} [\bar{\partial}_b, \zeta \Gamma_k \Psi^+ \zeta] v, w_{\zeta,k}^+)_{L^2} + (v_{\zeta,k}^+, [e^{2ks\sigma}, \bar{\partial}_b^*] w_{\zeta,k}^+)_{L^2} + (e^{2ks\sigma} v_{\zeta,k}^+, [\zeta \Gamma_k \Psi^+ \zeta, \bar{\partial}_b^*] w)_{L^2}| \\
&\leq \epsilon (\|e^{k(s\sigma+a)} [\bar{\partial}_b, \zeta \Gamma_k \Psi^+ \zeta] v\|_{L^2}^2 + \|e^{k(s\sigma+a)} v_{\zeta,+}^+\|_{L^2}^2) \\
&\quad + c_\epsilon (\|e^{k(s\sigma-a)} w_{\zeta,k}^+\|_{L^2}^2 + k^2 \|e^{k(s\sigma-a)} \partial_b \sigma \lrcorner w_{\zeta,k}^+\|_{L^2}^2 + \|e^{k(s\sigma-a)} [\zeta \Gamma_k \Psi^+ \zeta, \bar{\partial}_b^*] w\|_{L^2}^2). \tag{3.25}
\end{aligned}$$

Now multiply both sides (3.25) by $g^2(e^k)$ and sum over k . The desired estimate (3.23) follows by the estimates of commutator in Lemma 2.5. The desired estimate (3.24) is proved analogously. $\square$

Lemma 3.7. *Suppose that the σ -superlogarithmic property with the pair of rates (f, f_0) holds on $(0, q)$ -forms. Then the following holds:*

(i) *For $q \geq q_0$, we have*

$$\begin{aligned}
& \|u\|_{gf-\mathcal{A}_\zeta^+, s\sigma}^2 + \|u\|_{gf_0 \ln - \bar{\partial}_b \sigma \lrcorner \mathcal{A}_\zeta^+, s\sigma}^2 + \|\bar{\partial}_b^* u\|_{g-\mathcal{A}_\zeta^+, s\sigma}^2 \\
& \lesssim \text{Re}(\bar{\partial}_b \bar{\partial}_b^* u, u)_{g-\mathcal{A}_\zeta^+, s\sigma} + \|\bar{\partial}_b u\|_{g-\mathcal{A}_\zeta^+, s\sigma}^2 + \|u\|_{g-\mathcal{E}_{\zeta, \bar{\zeta}}^{s,1}}^2,
\end{aligned}$$

$$\text{for all } u \in (g-\mathcal{A}_{\zeta;0,q}^{+, s\sigma+1}) \cap (g-\mathcal{E}_{\zeta, \bar{\zeta};0,q}^{s,1}).$$

(ii) *For $q \leq n - q_0$, we have*

$$\begin{aligned}
& \|u\|_{gf-\mathcal{A}_\zeta^-, s\sigma}^2 + \|u\|_{gf_0 \ln - \bar{\partial}_b \sigma \wedge \mathcal{A}_\zeta^-, s\sigma}^2 + \|\bar{\partial}_b u\|_{g-\mathcal{A}_\zeta^-, s\sigma}^2 \\
& \lesssim \text{Re}(\bar{\partial}_b^* \bar{\partial}_b u, u)_{g-\mathcal{A}_\zeta^-, s\sigma} + \|\bar{\partial}_b u^*\|_{g-\mathcal{A}_\zeta^-, s\sigma}^2 + \|u\|_{g-\mathcal{E}_{\zeta, \bar{\zeta}}^{s,1}}^2,
\end{aligned}$$

$$\text{for all } u \in (g-\mathcal{A}_{\zeta;0,q}^{-, s\sigma+1}) \cap (g-\mathcal{E}_{\zeta, \bar{\zeta};0,q}^{s,1}) \text{ with } q \leq n - q_0.$$

Proof. Using Lemma 3.6 for $v = \bar{\partial}_b^* u$ and $w = u$ with the choice $a = 0$, we get

$$\begin{aligned}
& \|\bar{\partial}_b^* u\|_{g-\mathcal{A}_\zeta^+, s\sigma}^2 = \text{Re}(\bar{\partial}_b^* u, \bar{\partial}_b^* u)_{g-\mathcal{A}_\zeta^+, s\sigma} \\
& \leq \text{Re}(\bar{\partial}_b \bar{\partial}_b^* u, u)_{g-\mathcal{A}_\zeta^+, s\sigma} + \left| \text{Re}(\bar{\partial}_b^* u, \bar{\partial}_b^* u)_{g-\mathcal{A}_\zeta^+, s\sigma} - \text{Re}(\bar{\partial}_b \bar{\partial}_b^* u, u)_{g-\mathcal{A}_\zeta^+, s\sigma} \right| \\
& \leq \text{Re}(\bar{\partial}_b \bar{\partial}_b^* u, u)_{g-\mathcal{A}_\zeta^+, s\sigma} + \epsilon \left(\|\bar{\partial}_b^* u\|_{g-\mathcal{A}_\zeta^+, s\sigma}^2 + \|\bar{\partial}_b^* u\|_{g-\mathcal{E}_{\zeta, \bar{\zeta}}^{s,0}}^2 \right) \\
& \quad + c_\epsilon \left(\|u\|_{g-\mathcal{A}_\zeta^+, s\sigma}^2 + \|u\|_{g \ln - \bar{\partial}_b \sigma \lrcorner \mathcal{A}_\zeta^+, s\sigma}^2 + \|u\|_{g-\mathcal{E}_{\zeta, \bar{\zeta}}^{s,0}}^2 \right). \tag{3.26}
\end{aligned}$$

First, we see that for small $\epsilon > 0$, the term $\|\bar{\partial}_b^* u\|_{g-\mathcal{A}_\zeta^+, s\sigma}^2$ in the third line of (3.26) is absorbed. Then, adding $\|\bar{\partial}_b u\|_{g-\mathcal{A}_\zeta^+, s\sigma}^2$ to both sides of (3.26) and applying Theorem 3.1, we

absorb $c_\epsilon \left(\|u\|_{g-\mathcal{A}_\zeta^+, s\sigma}^2 + \|u\|_{g \ln - \partial_b \sigma \lrcorner \mathcal{A}_\zeta^+, s\sigma}^2 \right)$ by $\|u\|_{gf-\mathcal{A}_\zeta^+, s\sigma}^2 + \|u\|_{gf_0 \ln - \partial_b \sigma \lrcorner \mathcal{A}_\zeta^+, s\sigma}^2$ since $f, f_0 \gg 1$. Therefore, the remainder terms, after absorbing, are $\operatorname{Re}(\bar{\partial}_b \bar{\partial}_b^* u, u)_{g-\mathcal{A}_\zeta^+, s\sigma} + \|\bar{\partial}_b u\|_{g-\mathcal{A}_\zeta^+, s\sigma}^2$ and terms in the error norm $\|\cdot\|_{g-\mathcal{E}_{\zeta, \tilde{\zeta}}^{s,0}}^2$. Finally, we notice that all terms in the error norm are bounded by $\|u\|_{g-\mathcal{E}_{\zeta, \tilde{\zeta}}^{s,1}}^2$. This proves the first estimate in Lemma 3.6. The proof of the second follows analogously. $\square$

The main theorem in this subsection is the *a-priori* estimates for $\square_b$ on $\pm$ -microlocalizations of $(0, q)$ -forms in the norm $\|\cdot\|_{g-\mathcal{A}_\zeta^\pm, s\sigma}$.

Theorem 3.8. *Suppose that the σ -superlogarithmic property on $(0, q_0)$ -forms with the pair of rates (f, f_0) holds. Then, for $s, r \in \mathbb{R}^+$ with $r \geq s_0 + 2$, the following holds.*

(i) *For $u \in \mathcal{A}_{\zeta; 0, q}^{+, s\sigma+2-r} \cap \mathcal{E}_{\zeta, \tilde{\zeta}; 0, q}^{s, 2-r}$ with $q \geq q_0$, we have*

$$\begin{aligned} & \|u\|_{f^2-\mathcal{A}_\zeta^+, s\sigma-r}^2 + \|\bar{\partial}_b u\|_{f-\mathcal{A}_\zeta^+, s\sigma-r}^2 + \|\bar{\partial}_b^* u\|_{f-\mathcal{A}_\zeta^+, s\sigma-r}^2 + \|\bar{\partial}_b^* \bar{\partial}_b u\|_{\mathcal{A}_\zeta^+, s\sigma-r}^2 + \|\bar{\partial}_b \bar{\partial}_b^* u\|_{\mathcal{A}_\zeta^+, s\sigma-r}^2 \\ & \|u\|_{ff_0 \ln - \partial_b \sigma \lrcorner \mathcal{A}_\zeta^+, s\sigma-r}^2 + \|\bar{\partial}_b u\|_{f_0 \ln - \partial_b \sigma \lrcorner \mathcal{A}_\zeta^+, s\sigma-r}^2 \lesssim \|\square_b u\|_{\mathcal{A}_\zeta^+, s\sigma-r}^2 + \|u\|_{\mathcal{E}_{\zeta, \tilde{\zeta}}^{s, 2-r}}^2. \end{aligned}$$

(ii) *For $u \in \mathcal{A}_{\zeta; 0, q}^{-, s\sigma+2-r} \cap \mathcal{E}_{\zeta, \tilde{\zeta}; 0, q}^{s, 2-r}$ with $q \leq n - q_0$, we have*

$$\begin{aligned} & \|u\|_{f^2-\mathcal{A}_\zeta^-, s\sigma-r}^2 + \|\bar{\partial}_b u\|_{f-\mathcal{A}_\zeta^-, s\sigma-r}^2 + \|\bar{\partial}_b^* u\|_{f-\mathcal{A}_\zeta^-, s\sigma-r}^2 + \|\bar{\partial}_b^* \bar{\partial}_b u\|_{\mathcal{A}_\zeta^-, s\sigma-r}^2 + \|\bar{\partial}_b \bar{\partial}_b^* u\|_{\mathcal{A}_\zeta^-, s\sigma-r}^2 \\ & \|u\|_{ff_0 \ln - \bar{\partial}_b \sigma \wedge \mathcal{A}_\zeta^-, s\sigma-r}^2 + \|\bar{\partial}_b^* u\|_{f_0 \ln - \bar{\partial}_b \sigma \wedge \mathcal{A}_\zeta^-, s\sigma-r}^2 \lesssim \|\square_b u\|_{\mathcal{A}_\zeta^-, s\sigma-r}^2 + \|u\|_{\mathcal{E}_{\zeta, \tilde{\zeta}}^{s, 2-r}}^2. \end{aligned}$$

Proof. We only need to prove the positive microlocalization since the negative microlocalization follows analogously. Combining the estimate (3.1) in Theorem 3.1 with $\bar{\partial}_b u$ substituted in for “ u ” and Lemma 3.7(i) with the choice $g(t) = t^{-r} f(t)$ for some $r \geq s_0 + 2$, we get

$$\begin{aligned} & \|u\|_{f^2-\mathcal{A}_\zeta^+, s\sigma-r}^2 + \|\bar{\partial}_b^* u\|_{f-\mathcal{A}_\zeta^+, s\sigma-r}^2 + \|\bar{\partial}_b u\|_{f-\mathcal{A}_\zeta^+, s\sigma-r}^2 + \|\bar{\partial}_b u\|_{f_0 \ln - \partial_b \sigma \lrcorner \mathcal{A}_\zeta^+, s\sigma-r}^2 \\ & + \|u\|_{ff_0 \ln - \partial_b \sigma \lrcorner \mathcal{A}_\zeta^+, s\sigma-r}^2 + \|\bar{\partial}_b u\|_{f_0 \ln - \partial_b \sigma \lrcorner \mathcal{A}_\zeta^+, s\sigma-r}^2 \\ & \lesssim \operatorname{Re}(\bar{\partial}_b \bar{\partial}_b^* u, u)_{f-\mathcal{A}_\zeta^+, s\sigma-r} + \|\bar{\partial}_b^* \bar{\partial}_b u\|_{\mathcal{A}_\zeta^+, s\sigma-r}^2 + \|u\|_{\mathcal{E}_{\zeta, \tilde{\zeta}}^{s, 2-r}}^2. \end{aligned} \tag{3.27}$$

Using the Cauchy-Schwarz inequality,

$$\left| \operatorname{Re}(\bar{\partial}_b \bar{\partial}_b^* u, u)_{f-\mathcal{A}_\zeta^+, s\sigma-r} \right| \leq \epsilon \|u\|_{f^2-\mathcal{A}_\zeta^+, s\sigma-r}^2 + c_\epsilon \|\bar{\partial}_b \bar{\partial}_b^* u\|_{\mathcal{A}_\zeta^+, s\sigma-r}^2,$$

and absorbing $\|u\|_{f^2-\mathcal{A}_\zeta^+, s\sigma-r}^2$, we only need to bound $\|\bar{\partial}_b^* \bar{\partial}_b u\|_{\mathcal{A}_\zeta^+, s\sigma-r}^2 + \|\bar{\partial}_b \bar{\partial}_b^* u\|_{\mathcal{A}_\zeta^+, s\sigma-r}^2$. By the definition of $\square_b$, it is easy to see that

$$\|\bar{\partial}_b^* \bar{\partial}_b u\|_{\mathcal{A}_\zeta^+, s\sigma-r}^2 + \|\bar{\partial}_b \bar{\partial}_b^* u\|_{\mathcal{A}_\zeta^+, s\sigma-r}^2 = \|\square_b u\|_{\mathcal{A}_\zeta^+, s\sigma-r}^2 - 2\operatorname{Re}(\bar{\partial}_b \bar{\partial}_b^* u, \bar{\partial}_b^* \bar{\partial}_b u)_{\mathcal{A}_\zeta^+, s\sigma-r} \tag{3.28}$$

For the last term, we apply again Lemma 3.7(i) for $v = \bar{\partial}_b \bar{\partial}_b^* u$ and $w = \bar{\partial}_b u$ with the choice $g(t) = t^{-r}$ and $a = 0$, since $\bar{\partial}_b v = \bar{\partial}_b \bar{\partial}_b \bar{\partial}_b^* u = 0$ it follows

$$\begin{aligned} \left| \operatorname{Re}(\bar{\partial}_b \bar{\partial}_b^* u, \bar{\partial}_b^* \bar{\partial}_b u)_{\mathcal{A}_\zeta^{+, s\sigma-r}} \right| &\leq \epsilon \left(\|\bar{\partial}_b \bar{\partial}_b^* u\|_{\mathcal{A}_\zeta^{+, s\sigma-r}}^2 + \|\bar{\partial}_b \bar{\partial}_b^* u\|_{\mathcal{E}_{\zeta, \bar{\zeta}}^{s, -r}}^2 \right) \\ &\quad + c_\epsilon \left(\|\bar{\partial}_b u\|_{\mathcal{A}_\zeta^{+, s\sigma-r}}^2 + \|\bar{\partial}_b u\|_{\ln - \bar{\partial}_b \sigma \lrcorner \mathcal{A}_\zeta^{+, s\sigma-r}}^2 + \|\bar{\partial}_b u\|_{\mathcal{E}_{\zeta, \bar{\zeta}}^{s, -r}}^2 \right). \end{aligned} \quad (3.29)$$

From (3.27), (3.28), (3.29), and all terms in the error norms are bounded by $\|u\|_{\mathcal{E}_{\zeta, \bar{\zeta}}^{s, 2-r}}$, we obtain the desired inequality. It completes the proof of Theorem 3.8. $\square$

3.3. A-priori estimates for $\square_b^\delta$. For $\delta > 0$, we set

$$\square_b^\delta := \square_b + \delta (\square_d + I),$$

where $\square_d := d^* d + d d^*$ is the elliptic operator associated to the de Rham exterior derivative d . We will give more discussion on $\square_b^\delta$ in the next section. By the ellipticity of $\square_d$, we have the following equivalences

$$\begin{aligned} \|u\|_{\mathcal{A}_\zeta^{+, s\sigma+2-r}}^2 + \|u\|_{\mathcal{E}_{\zeta, \bar{\zeta}}^{s, 2-r}}^2 &\approx \|du\|_{\mathcal{A}_\zeta^{+, s\sigma+1-r}}^2 + \|d^* u\|_{\mathcal{A}_\zeta^{+, s\sigma+1-r}}^2 + \|u\|_{\mathcal{A}_\zeta^{+, s\sigma+1-r}}^2 + \|u\|_{\mathcal{E}_{\zeta, \bar{\zeta}}^{s, 2-r}}^2 \\ &\approx ((\square_d + I)u, u)_{\mathcal{A}_\zeta^{+, s\sigma+1-r}}^2 + \|u\|_{\mathcal{E}_{\zeta, \bar{\zeta}}^{s, 2-r}}^2 \\ &\approx \|((\square_d + I)u)\|_{\mathcal{A}_\zeta^{+, s\sigma-r}}^2 + \|u\|_{\mathcal{E}_{\zeta, \bar{\zeta}}^{s, 2-r}}^2 \end{aligned} \quad (3.30)$$

for any $u \in \mathcal{A}_{\zeta; 0, q}^{+, s\sigma+2-r} \cap \mathcal{E}_{\zeta, \bar{\zeta}; 0, q}^{s, 2-r}$ and $0 \leq q \leq n$. The boundedness of $\square_b$ by $\square_b^\delta$ is given in the following lemma.

Lemma 3.9. *Suppose that the σ -superlogarithmic property on $(0, q_0)$ -forms holds. Then the following holds uniformly in δ .*

(i) *For $u \in \mathcal{A}_{\zeta; 0, q}^{+, s\sigma+2-r} \cap \mathcal{E}_{\zeta, \bar{\zeta}; 0, q}^{s, 2-r}$ with $q \geq q_0$, we have*

$$\|\square_b u\|_{\mathcal{A}_\zeta^{+, s\sigma-r}}^2 \lesssim \|\square_b^\delta u\|_{\mathcal{A}_\zeta^{+, s\sigma-r}}^2 + \|\bar{\partial}_b u\|_{\mathcal{A}_\zeta^{+, s\sigma-r}}^2 + \|\bar{\partial}_b u\|_{\ln - \bar{\partial}_b \sigma \lrcorner \mathcal{A}_\zeta^{+, s\sigma-r}}^2 + \|u\|_{\mathcal{E}_{\zeta, \bar{\zeta}}^{s, 2-r}}^2.$$

ii. *For $u \in \mathcal{A}_{\zeta; 0, q}^{-, s\sigma+2-r} \cap \mathcal{E}_{\zeta, \bar{\zeta}; 0, q}^{s, 2-r}$ with $q \leq n - q_0$, we have*

$$\|\square_b u\|_{\mathcal{A}_\zeta^{-, s\sigma-r}}^2 \lesssim \|\square_b^\delta u\|_{\mathcal{A}_\zeta^{-, s\sigma-r}}^2 + \|\bar{\partial}_b^* u\|_{\mathcal{A}_\zeta^{-, s\sigma-r}}^2 + \|\bar{\partial}_b^* u\|_{\ln - \bar{\partial}_b \sigma \wedge \mathcal{A}_\zeta^{-, s\sigma-r}}^2 + \|u\|_{\mathcal{E}_{\zeta, \bar{\zeta}}^{s, 2-r}}^2.$$

Proof. Now, we use Lemma 3.7.(i) for the choice $g(t) = t^{-r+1}$ to get

$$\|\bar{\partial}_b^* u\|_{\mathcal{A}_\zeta^{+, s\sigma-r+1}}^2 \leq \operatorname{Re}(\bar{\partial}_b \bar{\partial}_b^* u, u)_{\mathcal{A}_\zeta^{+, s\sigma-r+1}} + \|\bar{\partial}_b u\|_{\mathcal{A}_\zeta^{+, s\sigma-r+1}}^2 + \|u\|_{\mathcal{E}_{\zeta, \bar{\zeta}}^{s, 2-r}}^2 \quad (3.31)$$

To estimate $\|\bar{\partial}_b u\|_{\mathcal{A}_\zeta^{+,s\sigma-r+1}}^2$, we apply Lemma 3.6.(i) for $v = u$ and $w = \bar{\partial}_b u$ with the choice $g(t) = t^{-r+1}$ and $a = 1$, $\epsilon := \epsilon\delta$ to obtain

$$\begin{aligned} \|\bar{\partial}_b u\|_{\mathcal{A}_\zeta^{+,s\sigma-r+1}}^2 &= \text{Re}(\bar{\partial}_b u, \bar{\partial}_b u)_{\mathcal{A}_\zeta^{+,s\sigma-r+1}} \\ &\leq \text{Re}(\bar{\partial}_b^* \bar{\partial}_b u, u)_{\mathcal{A}^{+,s\sigma-r+1}} + \epsilon\delta \left(\|u\|_{\mathcal{A}_\zeta^{+,s\sigma-r+2}}^2 + \|u\|_{\mathcal{E}_{\zeta,\tilde{\zeta}}^{s,2-r}}^2 \right) \\ &\quad + c_\epsilon \delta^{-1} \left(\|\bar{\partial}_b u\|_{\mathcal{A}_\zeta^{+,s\sigma-r}}^2 + \|\bar{\partial}_b u\|_{\ln-\partial_b \sigma \lrcorner \mathcal{A}_\zeta^{+,s\sigma-r}}^2 + \|\bar{\partial}_b u\|_{\mathcal{E}_\zeta^{s,-r}}^2 \right). \end{aligned} \quad (3.32)$$

Combining (3.31) and (3.32), we get

$$\begin{aligned} \|\bar{\partial}_b^* u\|_{\mathcal{A}_\zeta^{+,s\sigma-r+1}}^2 &\leq \text{Re}(\square_b u, u)_{\mathcal{A}_\zeta^{+,s\sigma-r+1}} + \epsilon\delta \|u\|_{\mathcal{A}_\zeta^{+,s\sigma-r+2}}^2 \\ &\quad + c_\epsilon \delta^{-1} \left(\|\bar{\partial}_b u\|_{\mathcal{A}_\zeta^{+,s\sigma-r}}^2 + \|\bar{\partial}_b u\|_{\ln-\partial_b \sigma \lrcorner \mathcal{A}_\zeta^{+,s\sigma-r}}^2 + \|u\|_{\mathcal{E}_{\zeta,\tilde{\zeta}}^{s,2-r}}^2 \right). \end{aligned} \quad (3.33)$$

Adding $(\delta(\square_d + I)u, u)_{\mathcal{A}_\zeta^{+,s\sigma+1-r}}$ into both sides of (3.33), after that using the Cauchy-Schwarz inequality for $\text{Re}(\square_b^\delta u, u)_{\mathcal{A}_\zeta^{+,s\sigma-r+1}}$, and then multiplying with δ , we obtain

$$\begin{aligned} \delta^2((\square_d + I)u, u)_{\mathcal{A}_\zeta^{+,s\sigma+1-r}} + \delta \|\bar{\partial}_b^* u\|_{\mathcal{A}_\zeta^{+,s\sigma-r+1}}^2 &\leq 2\epsilon\delta^2 \|u\|_{\mathcal{A}_\zeta^{+,s\sigma-r+2}}^2 \\ &\quad + c_\epsilon \left(\|\square_b^\delta u\|_{\mathcal{A}_\zeta^{+,s\sigma-r}}^2 + \|\bar{\partial}_b u\|_{\mathcal{A}_\zeta^{+,s\sigma-r}}^2 + \|\bar{\partial}_b u\|_{\ln-\partial_b \sigma \lrcorner \mathcal{A}_\zeta^{+,s\sigma-r}}^2 + \|u\|_{\mathcal{E}_{\zeta,\tilde{\zeta}}^{s,2-r}}^2 \right). \end{aligned} \quad (3.34)$$

Using (3.30) to absorb the first term in the second line of (3.34), we get the estimate for $\delta^2(\square_d + I)$ in the following inequality

$$\delta^2 \|(\square_d + I)u\|_{\mathcal{A}_\zeta^{+,s\sigma-r}}^2 \lesssim \|\square_b^\delta u\|_{\mathcal{A}_\zeta^{+,s\sigma-r}}^2 + \|\bar{\partial}_b u\|_{\mathcal{A}_\zeta^{+,s\sigma-r}}^2 + \|\bar{\partial}_b u\|_{\ln-\partial_b \sigma \lrcorner \mathcal{A}_\zeta^{+,s\sigma-r}}^2 + \|u\|_{\mathcal{E}_{\zeta,\tilde{\zeta}}^{s,2-r}}^2.$$

The proof of the first part of this lemma is complete by the simple inequality

$$\|\square_b u\|_{\mathcal{A}_\zeta^{+,s\sigma-r}}^2 \leq 2\delta^2 \|(\square_d + I)u\|_{\mathcal{A}_\zeta^{+,s\sigma-r}}^2 + \|\square_b^\delta u\|_{\mathcal{A}_\zeta^{+,s\sigma-r}}^2.$$

The proof of the second part goes in analogously way. $\square$

The estimates for elliptic regularization method follows immediately by Theorem 3.8 and Lemma 3.9.

Theorem 3.10. *Suppose that the σ -superlogarithmic property on $(0, q_0)$ -forms holds. Then, the following holds uniformly in δ .*

- (i) *For any $u \in \mathcal{A}_{\zeta;0,q}^{+,s\sigma+2-r} \cap \mathcal{E}_{\zeta,\tilde{\zeta};0,q}^{s,2-r}$ with $q \geq q_0$, we have*

$$\|u\|_{\mathcal{A}_\zeta^{+,s\sigma-r}}^2 \lesssim \|\square_b^\delta u\|_{\mathcal{A}_\zeta^{+,s\sigma-r}}^2 + \|u\|_{\mathcal{E}_{\zeta,\tilde{\zeta}}^{s,2-r}}^2.$$

ii. For any $u \in \mathcal{A}_{\zeta;0,q}^{-,s\sigma+2-r} \cap \mathcal{E}_{\zeta,\zeta;0,q}^{s,2-r}$ with $q \geq q_0$, we have

$$\|u\|_{\mathcal{A}_{\zeta}^{-,s\sigma-r}}^2 \lesssim \|\square_b^\delta u\|_{\mathcal{A}_{\zeta}^{+,s\sigma-r}}^2 + \|u\|_{\mathcal{E}_{\zeta,\zeta}^{s,2-r}}^2.$$

We finish this section by the elliptic estimate for 0-microlocalization.

Proposition 3.11. *Let $\Psi_0^0 \prec \Psi_1^0$ be a pair of homogeneous pseudodifferential operators of order zero whose symbols have support in $\mathcal{C}^0$ and let $\zeta \prec \zeta_1 \prec \zeta_2$ be a triple of cutoff functions. Then, for any $s, s_o \geq 0$ and $\delta \geq 0$, we have*

$$\|\Psi_0^0 \zeta u\|_{H^{s+1}}^2 \lesssim \|\Psi_1^0 \zeta_1 \bar{\partial}_b u\|_{H^s}^2 + \|\Psi_1^0 \zeta_1 \bar{\partial}_b^* u\|_{H^s}^2 + \|\zeta_2 u\|_{H^{-s_o}}^2, \quad (3.35)$$

$$\|\Psi_0^0 \zeta u\|_{H^{s+2}}^2 \lesssim \|\Psi_1^0 \zeta_1 \square_b u\|_{H^s}^2 + \|\zeta_2 u\|_{H^{-s_o}}^2, \quad (3.36)$$

$$\|\Psi_0^0 \zeta u\|_{H^{s+2}}^2 \lesssim \|\Psi_1^0 \zeta_1 \square_b^\delta u\|_{H^s}^2 + \|\zeta_2 u\|_{H^{-s_o}}^2, \quad (3.37)$$

for all $u \in C_{0,q}^\infty(\text{supp}(\zeta_1)) \cap H_{0,q}^{-s_o}(M)$ with any $0 \leq q \leq n$.

The proof of this proposition is omitted since it follows straightforwardly from the elliptic estimate for 0-microlocal.

4. PROOF OF THE MAIN THEOREMS

4.1. Local regularity of the complex Green operator. Let Q_b be defined by

$$Q_b(u, u) = \|\bar{\partial}_b u\|^2 + \|\bar{\partial}_b^* u\|^2, \quad u \in \text{Dom}(\bar{\partial}_b) \cap \text{Dom}(\bar{\partial}_b^*).$$

The closed range of $\bar{\partial}_b$ in L^2 -spaces for all degrees of forms implies that

$$\|u\|^2 \leq c (Q_b(u, u) + \|H_q u\|^2), \quad (4.1)$$

for all $(0, q)$ forms $u \in \text{Dom}(\bar{\partial}_b) \cap \text{Dom}(\bar{\partial}_b^*) \cap L_{0,q}^2(M)$ where H_q is the project onto $\mathcal{H}_{0,q}$ and $0 \leq q \leq n$. Here we recall that $\mathcal{H}_{0,q}$ is the space of harmonic forms of degree $(0, q)$. As a consequence of (4.1), $\square_b$ is invertible over $\mathcal{H}_{0,q}^\perp$ and its inverse G_q is L^2 bounded. as mentioned in Section 1, we extend the operator G_q to a linear operator on $L_{0,q}^2(M)$ by setting it equal to zero on $\mathcal{H}_{0,q}$. Then G_q is bounded and self adjoint in $L_{0,q}^2(M)$, and

$$\square_b G_q = I - H_q.$$

As a consequence from the L^2 boundedness of G_q , the related operators $\bar{\partial}_b^* G_q$, $G_q \bar{\partial}_b^*$, $\bar{\partial}_b G_q$, $G_q \bar{\partial}_b$, $I - \bar{\partial}_b^* \bar{\partial}_b G_q$, $I - \bar{\partial}_b^* G_q \bar{\partial}_b$, $I - \bar{\partial}_b \bar{\partial}_b^* G_q$, $I - \bar{\partial}_b G_q \bar{\partial}_b^*$, $\bar{\partial}_b G_q^2 \bar{\partial}_b^*$ and $\bar{\partial}_b^* G_q^2 \bar{\partial}_b$ are also L^2 bounded.

For higher regularity of G_q , by Theorem 3.8, Proposition 3.11 and the comparison of the $\|\cdot\|_{\mathcal{A}_{\zeta}^{s\sigma}}$ with $\|\zeta_0 \cdot\|_{H^s}$ and $\|\zeta_1 \cdot\|_{H^s}$ and the choice of cutoff functions $\zeta_0 \prec \sigma \prec \zeta \prec \tilde{\zeta} \prec \zeta_1 \prec \zeta_2$, we obtain the a-priori estimate in H^s : for any $s \geq 0$, there exists $c_s > 0$ so that for every $(0, q)$ form u smooth in U ,

$$\|\zeta_0 u\|_{H^s}^2 \leq c_s (\|\zeta_1 \square_b u\|_{H^s}^2 + \|u\|_{L^2}^2).$$

for all $u \in C_{0,q}^\infty(\text{supp}(\zeta_1)) \cap L_{0,q}^2(M)$ with $q_0 \leq q \leq n - q_0$. Unlike the L^2 -estimate (4.1), this does not imply $\zeta_0 G_q \varphi \in H_{0,q}^s(M)$ for $\varphi \in H_{0,q}^s(\text{supp}(\zeta_1)) \cap L_{0,q}^2(M)$ since we do not know whether $u = G_q \varphi \in \mathcal{A}_{\zeta;0,q}^{s\sigma}$ even that φ is smooth. We need to work with the family of regularized operators G_q^δ defined as follows.

For a small $\delta > 0$, we define

$$Q_b^\delta(u, v) := Q_b(u, v) + \delta Q_d(u, v) \quad u, v \in H_{0,q}^1(M),$$

where $Q_d(u, v) = (du, dv) + (d^*u, d^*v) + (u, v)$ the Hermitian inner product associated to the de Rham exterior derivative d . This quadratic form Q_b^δ gives a unique, self-adjoint in L^2 , elliptic operator $\square_b^\delta := \square_b + \delta(\square_d + I)$ which is mentioned in §3.3. Since $\delta\|u\|^2 \leq Q_b^\delta(u, u)$ for all $u \in H_{0,q}^1(M)$, $\square_b^\delta$ has an inverse G_q^δ such that

$$\square_b^\delta G_q^\delta = I \quad \text{since } \ker(\square_b^\delta) = \{0\}.$$

Furthermore, by the theory of elliptic regularity, it is well-known that G_q^δ is locally regular. More precisely, if $\varphi \in L_{0,q}^2(M) \cap H_{0,q}^s(V)$ then $G_q^\delta \varphi \in H_{0,q}^{s+2}(V)$ for any open set V of M . Consequently, G_q^δ is globally regular, i.e., its mapping is continuous in $C_{0,q}^\infty(M)$.

Let $H_{0,q}^{-\infty}(M) = \cup_{s \in \mathbb{R}} H_{0,q}^s(M)$ be the dual space of $C_{0,q}^\infty(M)$. Since G_q^δ is self-adjoint and its mapping is continuous in the $C_{0,q}^\infty(M)$ topology, it extends to a continuous in the space $H_{0,q}^{-\infty}(M)$. Thus, the local regularity property can be replaced as: if $\varphi \in H_{0,q}^{-\infty}(M) \cap H_{0,q}^s(V)$ then $G_q^\delta \varphi \in H_{0,q}^{s+2}(V)$ for any open set V of M . The following proposition is to show that $G_q \varphi \in \mathcal{A}_\zeta^{\pm, s\sigma-r}$ by the advantage of $G_q^\delta \varphi$.

Theorem 4.1. *Suppose that the σ -superlogarithmic property on $(0, q_0)$ -forms holds and $\sigma \prec \zeta \prec \tilde{\zeta} \prec \zeta_1 \prec \zeta_2$. Let $\varphi \in H_{0,q}^{-\infty}(M) \cap C_{0,q}^\infty(\text{supp}(\zeta_1))$ such that $\|\zeta_2 G_q^\delta \varphi\|_{H^{-s_0}}^2 < \infty$ uniformly in δ for some $s_0 \geq 0$. Then, for any $r \geq s_0 + 2$ and $s \geq 0$, we have*

- (i) $G_q \varphi \in \mathcal{A}_{\zeta;0,q}^{+, s\sigma-r}$ for $q_0 \leq q \leq n$;
- (ii) $G_q \varphi \in \mathcal{A}_{\zeta;0,q}^{-, s\sigma-r}$ for $0 \leq q \leq n - q_0$;
- (iii) $G_q \varphi \in \mathcal{E}_{\zeta, \tilde{\zeta};0,q}^{s, -r}$ for all $q = 0, \dots, n$.

Proof. By the local regularity of G_q^δ , we have $G_q^\delta \varphi \in C_{0,q}^\infty(\text{supp}(\zeta_1))$. Consequently, $G_q^\delta \varphi \in \mathcal{A}_{\zeta;0,q}^{+, s\sigma-r+2} \cap \mathcal{E}_{\zeta, \tilde{\zeta};0,q}^{s, -r+2}$ for all $s, r \geq 0$ with $r \geq s_0 + 2$. So we can apply Theorem 3.10(i) and Proposition 3.11(iii) for $u = G_q^\delta \varphi$ with $q \geq q_0$ to get the uniform estimate in δ

$$\begin{aligned} \|G_q^\delta \varphi\|_{\mathcal{A}_\zeta^{+, s\sigma-r}}^2 &\lesssim \|\varphi\|_{\mathcal{A}_\zeta^{+, s\sigma-r}}^2 + \|G_q^\delta \varphi\|_{\mathcal{E}_{\zeta, \tilde{\zeta}}^{s, -r+2}}^2 \\ &= \|\varphi\|_{\mathcal{A}_\zeta^{+, s\sigma-r}}^2 + \|\tilde{\Psi}^0 \zeta G_q^\delta \varphi\|_{H^{s-r+2}}^2 + \|\tilde{\zeta} G_q^\delta \varphi\|_{H^{-r+2}}^2 \\ &\lesssim \|\zeta_1 \varphi\|_{H^{s-r}}^2 + \|\zeta_2 G_q^\delta \varphi\|_{H^{-r+2}}^2. \end{aligned}$$

Thus the assumption that $\|\zeta_2 G_q^\delta \varphi\|_{H^{-r+2}}^2$ is bounded uniformly in δ forces $\|G_q^\delta \varphi\|_{\mathcal{A}_\zeta^{+,s\sigma-r}}^2$ to be also uniform bounded in δ . Consequently, the family $\{G_q^\delta \varphi\}_{0 < \delta \leq 1}$ has a subsequence which converges weakly to $\hat{u}$ in the norm $\|\cdot\|_{\mathcal{A}_\zeta^{+,s\sigma-r}}$. We claim $\hat{u} = G_q \varphi$ in $V_0^{\zeta_2}$ the interior of $\{x \in M : \zeta_2(x) = 1\}$. Let $Q_{b,V_0^{\zeta_2}}$ be the restrictions of Q_b to $V_0^{\zeta_2}$. For any $v \in C_{0,q}^\infty(M)$ supported in $V_0^{\zeta_2}$, we have

$$\begin{aligned} |Q_{b,V_0^{\zeta_2}}(G_q^\delta \varphi - G_q \varphi, v)| &= |Q_b(G_q^\delta \varphi - G_q \varphi, v)| = \delta |Q_d(G_q^\delta \varphi, v)| = \delta |Q_d(\zeta_2 G_q^\delta \varphi, v)| \\ &\lesssim \delta \|\zeta_2 G_q^\delta \varphi\|_{H^{2-r}} \|v\|_{H^r} \lesssim \delta \|v\|_{H^r} \rightarrow 0 \quad \text{as } \delta \rightarrow 0. \end{aligned}$$

It means that $G_q^\delta \varphi$ converges to $G_q \varphi$ weakly in the $Q_{b,V_0^{\zeta_2}}$ -norm. Therefore, we must have $\hat{u} = G_q \varphi$ in $V_0^{\zeta_2} \supset \text{supp}(\zeta)$ and hence

$$\|G_q \varphi\|_{\mathcal{A}_\zeta^{+,s\sigma-r}} \leq \liminf \|G_q^\delta \varphi\|_{\mathcal{A}_\zeta^{+,s\sigma-r}} < \infty.$$

Therefore, $G_q \varphi \in \mathcal{A}_{\zeta;0,q}^{+,s\sigma-r}$ for $q \geq q_0$. Similarly, $G_q \varphi \in \mathcal{A}_{\zeta;0,q}^{-,s\sigma-r}$ for $q \leq n - q_0$; and $G_q \varphi \in \mathcal{E}_{\zeta,\tilde{\zeta};0,q}^{s,-r}$ for all $0 \leq q \leq n$. $\square$

In the following theorem we give the proof of local hypoellipticity of $\square_b$ (the first part of Theorem 1.3).

Theorem 4.2. *Let M be a pseudoconvex CR manifold of dimension $(2n+1)$ with $n \geq 2$ such that $\bar{\partial}_b$ has closed range in L^2 spaces for all degrees of forms. Suppose that the σ -superlogarithmic property holds on $(0, q_0)$ -forms. Then, for all $q_0 \leq q \leq n - q_0$, if $u \in L_{0,q}^2(M)$ such that $\square_b u \in L_{0,q}^2(M) \cap C_{0,q}^\infty(V_1^\sigma)$ then $u \in C_{0,q}^\infty(V_0^\sigma)$.*

Proof. Set $\varphi := \square_b u \in \mathcal{H}_{0,q}^\perp(M) \cap L_{0,q}^2(M) \cap C_{0,q}^\infty(V_1^\sigma)$. Then $u = G_q \varphi + H_q u$ where we recall that H_q is the project onto $\mathcal{H}_{0,q}(M)$. In order to apply Theorem 3.10 above for this $G_q \varphi$, we have to show that $G_q^\delta \varphi$ is also bounded in $L_{0,q}^2(M)$ uniformly in δ . Since $\mathcal{H}_{0,q}$ is finite dimensional if $1 \leq q \leq n - 1$, the inequality 4.1 implies

$$\|u\|_{L^2}^2 \lesssim Q_b^\delta(u, u) + \|H_q^\delta u\|_{L^2}^2 \quad (4.2)$$

uniformly in δ where H_q^δ is the project onto $\mathcal{H}_{0,q}^\delta = \mathcal{H}_{0,q} \cap \{u : Q_d(u, u) = 0\} = \{0\}$ (see [KR, Lemma 5.3] for the proof of this claim). This implies

$$\|G_q^\delta \varphi\|_{L^2} \lesssim \|\varphi\|_{L^2} \quad (4.3)$$

holds uniformly in δ for any $\varphi \in L_{0,q}^2(M)$ with $1 \leq q \leq n - 1$. Thus, $\|\zeta_2 G_q^\delta \varphi\|_{H^0}^2 < \infty$ uniformly in δ . Now, we apply Theorem 4.1 for $s_o = 0$ and $r = 2$ to obtain that

- (i) $G_q \varphi \in \mathcal{A}_{\zeta;0,q}^{+,s\sigma-2}$ for $q_0 \leq q \leq n - 1$;
- (ii) $G_q \varphi \in \mathcal{A}_{\zeta;0,q}^{-,s\sigma-2}$ for $1 \leq q \leq n - q_0$;
- (iii) $G_q \varphi \in \mathcal{E}_{\zeta,\tilde{\zeta};0,q}^{s,-2}$ for $1 \leq q \leq n - 1$.

Subsisting $G_q\varphi \in \mathcal{A}_{\zeta;0,q}^{+,s\sigma-2} \cap \mathcal{A}_{\zeta;0,q}^{-,s\sigma-2} \cap \mathcal{E}_{\zeta,\tilde{\zeta};0,q}^{s,-2}$ with $q_0 \leq q \leq n - q_0$ into the estimates in Theorem 3.1, we get

$$\begin{aligned}
\|\zeta_0 u\|_{H^{s-4}}^2 &\lesssim \|\zeta_0 G_q \varphi\|_{H^{s-4}}^2 + \|\zeta_0 H_q u\|_{H^{s-4}}^2 \\
&\lesssim \|G_q \varphi\|_{\mathcal{A}_{\zeta}^{+,s\sigma-4}}^2 + \|G_q \varphi\|_{\mathcal{A}_{\zeta}^{-,s\sigma-4}}^2 + \|G_q \varphi\|_{\mathcal{E}_{\zeta,\tilde{\zeta}}^{s-4}}^2 + \|H_q u\|_{H^{s-4}}^2 \\
&\lesssim \|\varphi\|_{\mathcal{A}_{\zeta}^{+,s\sigma-4}}^2 + \|\varphi\|_{\mathcal{A}_{\zeta}^{-,s\sigma-4}}^2 + \|G_q \varphi\|_{\mathcal{E}_{\zeta,\tilde{\zeta}}^{s-4}}^2 + \|u\|_{L^2}^2 \\
&\lesssim \|\zeta_1 \varphi\|_{H^{s-4}}^2 + \|G_q \varphi\|_{L^2}^2 + \|u\|_{L^2}^2 \\
&\lesssim \|\zeta_1 \square_b u\|_{H^{s-4}}^2 + \|u\|_{L^2}^2.
\end{aligned} \tag{4.4}$$

Here, we have use $\|H_q u\|_{H^{s-4}}^2 \lesssim \|u\|_{L^2}^2$ since $\mathcal{H}_{0,q}(M)$ is finite dimensional. We conclude that $\zeta_0 u \in H^{s-4}$ for any $s \geq 0$ and any $\zeta_0 \in V_0^\sigma$, this proves Theorem 4.2. $\square$

Remark 4.3. As in the proof of this theorem, we see that even *a-priori* estimates hold on the top degrees, we only have the regularity of G_q excluding the top degrees $q = 0$ and $q = n$ since $\mathcal{H}_{0,q}$ is finite dimensional only for $1 \leq q \leq n - 1$.

The local regularity of G_q is equivalent to the local hypoellipticity of $\square_b$ and implies the local regularity of $\bar{\partial}_b G_q$ and other operators but with loss derivatives in H^s -spaces. In the following we prove the exact H^s regularity or the H^s regularity with a small gain derivative for the relative operators of G_q .

Theorem 4.4. *Let M be a pseudoconvex CR manifold of dimension $(2n + 1)$, $n \geq 2$ such that $\bar{\partial}_b$ has closed range in L^2 spaces for all degrees of forms. Suppose that the σ -superlogarithmic property with independent rate f holds on $(0, q_0)$ -forms. Then, for all $s \geq 0$, $q_0 \leq q \leq n - q_0$ and $\zeta_0 \prec \sigma \prec \zeta_1$, the following holds.*

(i) *If $\varphi \in L_{0,q}^2(M) \cap H_{0,q}^s(\text{supp}(\zeta_1))$, then*

$$\begin{aligned}
&\|f^2(\Lambda)\zeta_0 G_q \varphi\|_{H^s} + \|f(\Lambda)\zeta_0 \bar{\partial}_b G_q \varphi\|_{H^s} + \|f(\Lambda)\zeta_0 \bar{\partial}_b^* G_q \varphi\|_{H^s} \\
&\quad + \|\zeta_0 \bar{\partial}_b \bar{\partial}_b^* G_q \varphi\|_{H^s} + \|\zeta_0 \bar{\partial}_b^* \bar{\partial}_b G_q \varphi\|_{H^s} \lesssim \|\zeta_1 \varphi\|_{H^s} + \|\varphi\|_{L^2}.
\end{aligned} \tag{4.5}$$

(ii) *If $\varphi \in L_{q-1}^2(M) \cap H_{0,q-1}^s(\text{supp}(\zeta_1))$ then*

$$\begin{aligned}
&\|f^2(\Lambda)\zeta_0 \bar{\partial}_b^* G_q^2 \bar{\partial}_b \varphi\|_{H^s} + \|f(\Lambda)\zeta_0 G_q \bar{\partial}_b \varphi\|_{H^s} + \|\zeta_0 (I - \bar{\partial}_b^* G_q \bar{\partial}_b) \varphi\|_{H^s} \\
&\quad \|\zeta_1 \varphi\|_{H^s} + \|\varphi\|_{L^2}.
\end{aligned} \tag{4.6}$$

(iii) *If $\varphi \in L_{0,q+1}^2(M) \cap H_{0,q+1}^s(\text{supp}(\zeta_1))$*

$$\begin{aligned}
&\|f^2(\Lambda)\zeta_0 \bar{\partial}_b G_q^2 \bar{\partial}_b^* \varphi\|_{H^s} + \|f(\Lambda)\zeta_0 G_q \bar{\partial}_b^* \varphi\|_{H^s} + \|\zeta_0 (I - \bar{\partial}_b G_q \bar{\partial}_b^*) \varphi\|_{H^s} \\
&\quad \|\zeta_1 \varphi\|_{H^s} + \|\varphi\|_{L^2}.
\end{aligned} \tag{4.7}$$

Proof. In the proof of Theorem 4.2 we have already proven that $G_q \varphi \in \mathcal{A}_{\zeta;0,q}^{+,s\sigma-2}$ with $q_0 \leq q \leq n - 1$, $G_q \varphi \in \mathcal{A}_{\zeta;0,q}^{-,s\sigma-2}$ with $1 \leq q \leq n - q_0$, and also $G_q \varphi \in \mathcal{E}_{\zeta,\tilde{\zeta};0,q}^{s,-2}$ for

$1 \leq q \leq n-1$ if provided $\varphi \in L_{0,q}^2(M) \cap H_{0,q}^s(\text{supp}(\zeta_1))$. Hence, $G_q\varphi$ satisfies the hypothesis of Theorem 3.1 for $r = 4$, therefore, the estimates

$$\begin{aligned} & \|G_q\varphi\|_{f^2-\mathcal{A}_\zeta^+, s\sigma-4}^2 + \|\bar{\partial}_b G_q\varphi\|_{f-\mathcal{A}_\zeta^+, s\sigma-4}^2 + \|\bar{\partial}_b^* G_q\varphi\|_{f-\mathcal{A}_\zeta^+, s\sigma-4}^2 \\ & + \|\bar{\partial}_b^* \bar{\partial}_b G_q\varphi\|_{\mathcal{A}_\zeta^+, s\sigma-4}^2 + \|\bar{\partial}_b \bar{\partial}_b^* G_q\varphi\|_{\mathcal{A}_\zeta^+, s\sigma-4}^2 \lesssim \|\zeta_1\varphi\|_{H^{s-4}}^2 + \|\varphi\|_{L^2}^2 \end{aligned} \quad (4.8)$$

holds on degrees $q_0 \leq q \leq n-1$; and dually,

$$\begin{aligned} & \|G_q\varphi\|_{f^2-\mathcal{A}_\zeta^-, s\sigma-4}^2 + \|\bar{\partial}_b G_q\varphi\|_{f-\mathcal{A}_\zeta^-, s\sigma-4}^2 + \|\bar{\partial}_b^* G_q\varphi\|_{f-\mathcal{A}_\zeta^-, s\sigma-4}^2 \\ & + \|\bar{\partial}_b^* \bar{\partial}_b G_q\varphi\|_{\mathcal{A}_\zeta^-, s\sigma-4}^2 + \|\bar{\partial}_b \bar{\partial}_b^* G_q\varphi\|_{\mathcal{A}_\zeta^-, s\sigma-4}^2 \lesssim \|\zeta_1\varphi\|_{H^{s-4}}^2 + \|\varphi\|_{L^2}^2 \end{aligned} \quad (4.9)$$

holds on degrees $1 \leq q \leq n - q_0$.

Proof of (i). Fix $q_0 \leq q \leq n - q_0$. If $\varphi \in L_{0,q}^2(M) \cap H_{0,q}^s(\text{supp}(\zeta_1))$, then both (4.8) and (4.9) hold, so taking summation to get

$$\begin{aligned} & \|f^2(\Lambda)\zeta_0 G_q\varphi\|_{H^{s-4}}^2 + \|f(\Lambda)\zeta_0 \bar{\partial}_b G_q\varphi\|_{H^{s-4}}^2 + \|f(\Lambda)\zeta_0 \bar{\partial}_b^* G_q\varphi\|_{H^{s-4}}^2 \\ & + \|\zeta_0 \bar{\partial}_b^* \bar{\partial}_b G_q\varphi\|_{H^{s-4}}^2 + \|\zeta_0 \bar{\partial}_b \bar{\partial}_b^* G_q\varphi\|_{H^{s-4}}^2 \lesssim \|\zeta_1\varphi\|_{H^{s-4}}^2 + \|\varphi\|_{L^2}^2. \end{aligned} \quad (4.10)$$

Since $H_{0,q}^s(M)$ is dense in $H_{0,q}^{s-4}(M)$, the estimate (4.10) is still holds for $\varphi \in L_{0,q}^2(M) \cap H_{0,q}^{s-4}(\text{supp}(\zeta_1))$. This proves the first estimate in Theorem 4.4.

Proof of (ii). Now we deal $\varphi \in L_{q-1}^2(M) \cap H_{0,q-1}^s(\text{supp}(\zeta_1))$. By (4.2), $\|G_q^\delta \bar{\partial}_b \varphi\|_{L^2}$ is uniformly bounded and hence, by Theorem 4.2, $G_q \bar{\partial}_b \varphi \in \mathcal{A}_{\zeta;0,q}^{\pm, s\sigma-2} \cap \mathcal{E}_{\zeta,\bar{\zeta};0,q}^{s,-2}$. We first give estimate on the positive norm $\|\cdot\|_{\mathcal{A}_\zeta^+, s\sigma-4}$. We use Lemma 3.7.(i) for $u = G_q \bar{\partial}_b \varphi$ and with the choice $g(t) = t^{-4}$,

$$\begin{aligned} & \|G_q \bar{\partial}_b \varphi\|_{f-\mathcal{A}_\zeta^+, s\sigma-4}^2 + \|G_q \bar{\partial}_b \varphi\|_{f_0 \ln - \partial_b \sigma \lrcorner \mathcal{A}_\zeta^+, s\sigma-4}^2 + \|\bar{\partial}_b^* G_q \bar{\partial}_b \varphi\|_{\mathcal{A}_\zeta^+, s\sigma-4}^2 \\ & \lesssim \text{Re}(\bar{\partial}_b \bar{\partial}_b^* G_q \bar{\partial}_b \varphi, G_q \bar{\partial}_b \varphi)_{\mathcal{A}_\zeta^+, s\sigma-4} + \|\bar{\partial}_b G_q \bar{\partial}_b \varphi\|_{\mathcal{A}_\zeta^+, s\sigma-4}^2 + \|G_q \bar{\partial}_b \varphi\|_{\mathcal{E}_{\zeta,\bar{\zeta}}^{s,-4}}^2 \\ & = \text{Re}(\bar{\partial}_b \varphi, G_q \bar{\partial}_b \varphi)_{\mathcal{A}_\zeta^+, s\sigma-4} + \|\varphi\|_{\mathcal{E}^{s,-4}}^2 + \|G_q \bar{\partial}_b \varphi\|_{\mathcal{E}^{s,-3}}^2. \end{aligned} \quad (4.11)$$

Using Lemma 3.6.(i) for $v = \varphi$, $w = G_q \bar{\partial}_b \varphi$ with the choice $g = t^{-4}$ and $a = \epsilon = 1$

$$\begin{aligned} & \text{Re}(\bar{\partial}_b \varphi, G_q \bar{\partial}_b \varphi)_{\mathcal{A}_\zeta^+, s\sigma-4} \leq \text{Re}(\varphi, \bar{\partial}_b^* G_q \bar{\partial}_b \varphi)_{\mathcal{A}_\zeta^+, s\sigma-4} + \|\varphi\|_{\mathcal{A}_\zeta^+, s\sigma-4} + \|\varphi\|_{\mathcal{E}_{\zeta,\bar{\zeta}}^{s,-4}} \\ & + c \left(\|G_q \bar{\partial}_b \varphi\|_{\mathcal{A}_\zeta^+, s\sigma-4}^2 + \|G_q \bar{\partial}_b \varphi\|_{\ln - \partial_b \sigma \lrcorner \mathcal{A}_\zeta^+, s\sigma-4}^2 + \|G_q \bar{\partial}_b \varphi\|_{\mathcal{E}_{\zeta,\bar{\zeta}}^{s,-4}}^2 \right) \\ & \leq \epsilon \|\bar{\partial}_b^* G_q \bar{\partial}_b \varphi\|_{\mathcal{A}_\zeta^+, s\sigma-4}^2 + c_\epsilon \|\varphi\|_{\mathcal{A}_\zeta^+, s\sigma-4} + \|\varphi\|_{\mathcal{E}_{\zeta,\bar{\zeta}}^{s,-4}} \\ & + c \left(\|G_q \bar{\partial}_b \varphi\|_{\mathcal{A}_\zeta^+, s\sigma-4}^2 + \|G_q \bar{\partial}_b \varphi\|_{\ln - \partial_b \sigma \lrcorner \mathcal{A}_\zeta^+, s\sigma-4}^2 + \|G_q \bar{\partial}_b \varphi\|_{\mathcal{E}_{\zeta,\bar{\zeta}}^{s,-4}}^2 \right). \end{aligned} \quad (4.12)$$

Combining (4.11) and (4.12) and the elliptic estimate in the error norm, we conclude that

$$\|G_q \bar{\partial}_b \varphi\|_{f-\mathcal{A}_\zeta^+, s\sigma-4}^2 + \|\bar{\partial}_b^* G_{q_0} \bar{\partial}_b \varphi\|_{\mathcal{A}_\zeta^+, s\sigma-4}^2 \lesssim \|\zeta_1 \varphi\|_{H^{s-4}}^2 + \|\varphi\|_{L^2}^2. \quad (4.13)$$

Now we give estimate on the negative norm $\|\cdot\|_{\mathcal{A}_\zeta^-, s\sigma-4}$. We consider two case of q . If $1 \leq q-1 \leq n-q_0$, we use the fact that $G_q \bar{\partial}_b = \bar{\partial}_b G_{q-1}$ and (4.9) holds for all degree between 1 and $n-q_0$, then

$$\|G_q \bar{\partial}_b \varphi\|_{f-\mathcal{A}_\zeta^-, s\sigma-4}^2 + \|(I - \bar{\partial}_b^* G_q \bar{\partial}_b) \varphi\|_{\mathcal{A}_\zeta^-, s\sigma-4}^2 \lesssim \|\zeta_1 \varphi\|_{H^{s-4}}^2 + \|\varphi\|_{L^2}^2. \quad (4.14)$$

Otherwise, $q = 1$, we give a directly proof of (4.14) as the following

$$\begin{aligned} \|G_1 \bar{\partial}_b \varphi\|_{f-\mathcal{A}_\zeta^-, s\sigma-4}^2 &\lesssim \|\bar{\partial}_b^* G_1 \bar{\partial}_b \varphi\|_{\mathcal{A}_\zeta^-, s\sigma-4}^2 + \|G_1 \bar{\partial}_b \varphi\|_{\mathcal{E}_{\zeta, \bar{\zeta}}^{s, -4}}^2 \\ &\lesssim \|(I - \bar{\partial}_b^* G_1 \bar{\partial}_b) \varphi\|_{\mathcal{A}_\zeta^-, s\sigma-4}^2 + \|\varphi\|_{\mathcal{A}_\zeta^-, s\sigma-4}^2 + \|G_1 \bar{\partial}_b \varphi\|_{\mathcal{E}_{\zeta, \bar{\zeta}}^{s, -4}}^2 \\ &\lesssim \|\varphi\|_{\mathcal{A}_\zeta^-, s\sigma-4}^2 + \|(I - \bar{\partial}_b^* G_1 \bar{\partial}_b) \varphi\|_{\mathcal{E}_{\zeta, \bar{\zeta}}^{s, -4}}^2 + \|G_1 \bar{\partial}_b \varphi\|_{\mathcal{E}_{\zeta, \bar{\zeta}}^{s, -4}}^2. \end{aligned} \quad (4.15)$$

Here, we have used Theorem 3.1.(ii) twice: the first one for the first inequality and the second one for the third inequality with notice that $\bar{\partial}_b(I - \bar{\partial}_b^* G_1 \bar{\partial}_b) \varphi = \bar{\partial}_b^*(I - \bar{\partial}_b^* G_1 \bar{\partial}_b) \varphi = 0$. So (4.14) holds for both case of $q \leq n - q_0$.

Finally, the estimate of the term $\bar{\partial}_b^* G_q^2 \bar{\partial}_b \varphi$ follows by the estimates of $\bar{\partial}_b G_q$ and $G_q \bar{\partial}_b$. Summarizing, we get

$$\begin{aligned} \|f^2(\Lambda) \zeta_0 \bar{\partial}_b^* G_q^2 \bar{\partial}_b \varphi\|_{H^{s-4}} + \|f(\Lambda) \zeta_0 G_q \bar{\partial}_b \varphi\|_{H^{s-4}} + \|\zeta_0 (I - \bar{\partial}_b^* G_q \bar{\partial}_b) \varphi\|_{H^{s-4}} \\ \|\zeta_1 \varphi\|_{H^{s-4}} + \|\varphi\|_{L^2}, \end{aligned} \quad (4.16)$$

for all $\varphi \in L_{0, q-1}^2(M) \cap H_{0, q-1}^s(\text{supp}(\zeta_1))$. By the density of $H_{0, q}^s(M)$ in $H_{0, q}^{s-4}(M)$, we can replace the assumption $\varphi \in L_{0, q-1}^2(M) \cap H_{0, q-1}^s(\text{supp}(\zeta_1))$ by $\varphi \in L_{0, q-1}^2(M) \cap H_{0, q-1}^{s-4}(\zeta_1)$. This proves the part (ii) in Theorem (4.4). The part (iii) follows analogously. $\square$

Remark 4.5. It is known that on the top degrees, the Green operators G_0 and G_n agree with $\bar{\partial}_b^* G_1^2 \bar{\partial}_b$ and $\bar{\partial}_b G_{n-1}^2 \bar{\partial}_b^*$, respectively; and also $\bar{\partial}_b G_0 = G_1 \bar{\partial}_b$, $\bar{\partial}_b^* G_n = G_{n-1} \bar{\partial}_b^*$. Therefore, if $q_0 = 1$ then

$$\|f^2(\Lambda) \zeta_0 G_0 \varphi\|_{H^s} + \|f(\Lambda) \zeta_0 \bar{\partial}_b G_0 \varphi\|_{H^s} \lesssim \|\zeta_1 \varphi\|_{H^s} + \|\varphi\|_{L^2} \quad (4.17)$$

holds for all $\varphi \in L^2(M) \cap H^s(\text{supp}(\zeta_1))$; and

$$\|f^2(\Lambda) \zeta_0 G_n \varphi\|_{H^s} + \|f(\Lambda) \zeta_0 \bar{\partial}_b^* G_n \varphi\|_{H^s} \lesssim \|\zeta_1 \varphi\|_{H^s} + \|\varphi\|_{L^2} \quad (4.18)$$

holds for all $\varphi \in L_{0, n}^2(M) \cap H_{0, n}^s(\text{supp}(\zeta_1))$.

For the case $n \geq 2$, the local regularity of G_0 (resp. G_n) is inherited from the local regularity of G_1 (resp. G_{n-1}). In the case $n = 1$, this procedure does not work. In this case, in order to use a-priori estimates, we assume that G_0 (and hence G_1) is globally regular.

Theorem 4.6. *Let M be a 3-dimensional pseudoconvex CR manifold such that $\bar{\partial}_b$ has closed range on functions and G_0 is globally regular. Suppose that σ -superlogarithmic property with dependent rate f holds on $(0, 1)$ -forms. Then, for $q = 0, 1$, if $u \in L^2_{0,q}(M)$ such that $\square_b u \in L^2_{0,q}(M) \cap C^\infty_{0,q}(V_1^\sigma)$ then $u \in C^\infty_{0,q}(V_0^\sigma)$. Furthermore,*

$$\|f^2(\Lambda)\zeta_0 G_0 \varphi\|_{H^s} + \|f(\Lambda)\zeta_0 \bar{\partial}_b G_0 \varphi\|_{H^s} + \|\zeta_0(I - \bar{\partial}_b^* G_1 \bar{\partial}_b)\varphi\|_{H^s} \lesssim \|\zeta_1 \varphi\|_{H^s} + \|\varphi\|_{L^2} \quad (4.19)$$

holds for all $\varphi \in L^2(M) \cap H^s(\text{supp}(\zeta_1))$, and

$$\|f^2(\Lambda)\zeta_0 G_1 \varphi\|_{H^s} + \|f(\Lambda)\zeta_0 \bar{\partial}_b^* G_1 \varphi\|_{H^s} + \|\zeta_0(I - \bar{\partial}_b G_0 \bar{\partial}_b^*)\varphi\|_{H^s} \lesssim \|\zeta_1 \varphi\|_{H^s} + \|\varphi\|_{L^2} \quad (4.20)$$

holds for all $\varphi \in L^2_{0,1}(M) \cap H^s_{0,1}(\text{supp}(\zeta_1))$.

The idea of the proof of this theorem follows by the argument in [Str10, page 70].

Proof. We remark that the global regularity property of G_0 implies that $\bar{\partial}_b G_0$, $(I - \bar{\partial}_b G_0 \bar{\partial}_b^*)$, G_1 , $\bar{\partial}_b^* G_1$ and $(I - \bar{\partial}_b^* G_1 \bar{\partial}_b)$ are also global regularity. Thus, for $\varphi \in C^\infty(M)$, we can use the *a-priori* estimates in Theorem 3.1 and 3.8, and follow the proof in Theorem 4.4(ii) (for the case $n \geq 2$, but it still hold for $n = 1$) together with the closed range of $\bar{\partial}_b$ to get (4.19) and (4.20) for this globally smooth function φ . So we have to pass the smoothness of φ from global to local. In the following we only give the proof for G_0 since other operators follows analogously. By the fact that the local hypoellipticity of $\square_b$ on functions follows by the local regularity of G_0 , we have obtained

$$\|f^2(\Lambda)\zeta_0 G_0 \varphi\|_{H^s}^2 \lesssim \|\tilde{\zeta} \varphi\|_{H^s}^2 + \|\varphi\|_{L^2}^2 \quad (4.21)$$

for all $\varphi \in C^\infty(M)$. Here we replace ζ_1 by $\tilde{\zeta}$ such that $\sigma \prec \tilde{\zeta} \prec \zeta_1$ since ζ_1 can be chosen arbitrary (but dominating σ).

For a general $\varphi \in H^s(\text{supp}(\zeta_1)) \cap L^2(M)$, we choose a sequence of functions $\varphi_m \in C^\infty(M)$ such that $\varphi_m \rightarrow \varphi$ in $L^2(M)$. Since $\text{supp}(\zeta_1)$ and $\text{supp}((1 - \zeta_2))$ are disjoint, we use (4.21) for the smooth datum function $(1 - \zeta_2)(\varphi_l - \varphi_m)$ and get

$$\|f^2(\Lambda)\zeta_0 G_0(1 - \zeta_2)(\varphi_l - \varphi_m)\|_{H^s}^2 \lesssim \|\varphi_l - \varphi_m\|^2.$$

Consequently, the sequence $\{f^2(\Lambda)\zeta_0 G_0(1 - \zeta_2)\varphi_m\}$ is Cauchy in $H^s(M)$. Since it converges to $f^2(\Lambda)\zeta_0 G_0(1 - \zeta_2)\varphi$ in $L^2(M)$, thus $f^2(\Lambda)\zeta_0 G_0(1 - \zeta_2)\varphi \in H^s(M)$ and

$$\|f^2(\Lambda)\zeta_0 G_0(1 - \zeta_2)\varphi\|_{H^s}^2 \lesssim \|\varphi\|^2. \quad (4.22)$$

Since $\zeta_1 \varphi \in C^\infty(M)$, we used (4.21) for the datum $\zeta_1 \varphi$,

$$\|f^2(\Lambda)\zeta_0 G_0 \zeta_1 \varphi\|_{H^s}^2 \lesssim \|\zeta_1 \tilde{\zeta} \varphi\|_{H^s}^2 + \|\zeta_1 \varphi\|_{L^2}^2 \lesssim \|\zeta_1 \varphi\|_{H^s}^2 + \|\varphi\|_{L^2}^2. \quad (4.23)$$

Combining (4.22) and (4.23), we get the desired estimate:

$$\|\zeta_0 G_0 \varphi\|_{H^s}^2 \lesssim \|f^2(\Lambda)\zeta_0 G_0 \zeta_1 \varphi\|_{H^s}^2 + \|f^2(\Lambda)\zeta_0 G_0(1 - \zeta_2)\varphi\|_{H^s}^2 \lesssim \|\zeta_1 \varphi\|_{H^s}^2 + \|\varphi\|_{L^2}^2,$$

for all $\varphi \in H^s(\text{supp}(\zeta_1)) \cap L^2(M)$.

□

4.2. Smoothness of the complex Green kernel. Let $\mathcal{G}_q(\hat{x}, x)$, $0 \leq q \leq n$, be the integral kernel of the complex Green operator G_q , it means, G_q has the integral representation

$$G_q \varphi(x) = \int_M \langle \varphi(\hat{x}), \mathcal{G}_q(\hat{x}, x) dV(\hat{x}) = (\varphi(\bullet), \mathcal{G}_q(\bullet, x))_{L^2}$$

provided the integral exists; this imposes regularity conditions on φ and $\mathcal{G}_q$. Thus, $\mathcal{G}_q$ is a double distribution form of degree $(0, q; q, 0)$ on the product manifold $M \times M$. $\mathcal{G}_q$ can be expressed in local coordinate $\hat{U} \times U$ as

$$\mathcal{G}_q(\hat{x}, x) = \sum_{\hat{I}, I \in \mathcal{I}_q} \mathcal{G}_{\hat{I}, I}(\hat{x}, x) \bar{\omega}_{\hat{I}}(\hat{x}) \wedge \omega_I(x).$$

To show the smoothness of $\mathcal{G}_q$, we will prove that functions $\mathcal{G}_{\hat{I}, I}$ are smooth for all $\hat{I}, I \in \mathcal{I}_q$. Fix $\hat{I} \in \mathcal{I}_q$. Let $\delta_{\hat{x}_0}$ be the delta Dirac “function” with support in $\hat{x}_0 \in M$. We choose a $(0, q)$ -form datum

$$\varphi_{\hat{x}_0}(\hat{x}) = (D^{\hat{\alpha}})^* \delta_{\hat{x}_0}(\hat{x}) \bar{\omega}_{\hat{I}}(\hat{x}),$$

where $(D^{\hat{\alpha}})^*$ is the L^2 adjoint of the differential operator $D^{\hat{\alpha}}$. This data is smooth on $M \setminus \{\hat{x}_0\}$ and bounded in $H^{-(n+1+|\hat{\alpha}|)}$ -norm since

$$\begin{aligned} \|\varphi_{\hat{x}_0}\|_{H^{-(n+1+|\hat{\alpha}|)}}^2 &\lesssim \|\delta_{\hat{x}_0}\|_{H^{-(n+1)}}^2 \\ &= \int_{\mathbb{R}^{2n+1}} (1 + |\xi|^2)^{-(n+1)} \mathcal{F}\{\delta_{\hat{x}_0}\}(\xi) d\xi \\ &= \int_{\mathbb{R}^{2n+1}} (1 + |\xi|^2)^{-(n+1)} d\xi \lesssim 1. \end{aligned} \tag{4.24}$$

Thus,

$$\begin{aligned} G_q \varphi_{\hat{x}_0}(x) &= (\varphi_{\hat{x}_0}(\bullet), \mathcal{G}_q(\bullet, x))_{L^2} \\ &= \sum_{I \in \mathcal{I}_q} \left((D^{\hat{\alpha}})^* \delta_{\hat{x}_0}(\bullet), \mathcal{G}_{\hat{I}, I}(\bullet, x) \right)_{L^2} \bar{\omega}_I(x) \\ &= \sum_{I \in \mathcal{I}_q} \left(\delta_{\hat{x}_0}(\bullet), D^{\hat{\alpha}} \mathcal{G}_{\hat{I}, I}(\bullet, x) \right)_{L^2} \bar{\omega}_I(x) \\ &= \sum_{I \in \mathcal{I}_q} \overline{D_{\hat{x}_0}^{\hat{\alpha}} \mathcal{G}_{\hat{I}, I}(\hat{x}_0, x)} \bar{\omega}_I(x), \end{aligned} \tag{4.25}$$

and consequently,

$$\sum_{I \in \mathcal{I}_q} \left| D_x^{\alpha} D_{\hat{x}_0}^{\hat{\alpha}} \mathcal{G}_{\hat{I}, I}(\hat{x}_0, x) \right|^2 = \left| \overline{D_x^{\alpha}} G_q \varphi_{\hat{x}_0}(x) \right|^2. \tag{4.26}$$

Let ζ_0 be a cutoff function in U , we choose $x_0 \in V_0^{\zeta_0}$ then by the Sobolev lemma

$$\left| D_{x_0}^\alpha D_{\hat{x}_0}^{\hat{\alpha}} \mathcal{G}_{\hat{I}, I}(\hat{x}_0, x_0) \right| \leq \sup_{x \in M} \left| \overline{D_x^\alpha} \zeta_0 G_q \varphi_{\hat{x}_0}(x) \right| \lesssim \|\zeta_0 G_q \varphi_{\hat{x}_0}\|_{n+1+|\alpha|}, \quad (4.27)$$

for all $I, \hat{I} \in \mathcal{I}_q$. These equalities (4.25), (4.26) and inequalities (4.27) hold if provided that $\|\zeta_0 G_q \varphi_{\hat{x}_0}\|_{H^{n+1+|\alpha|}} < \infty$. To show this boundedness, we need the hypothesis of weakly superlogarithmic properties. Theorem 1.6 is splitted into two following theorems.

Theorem 4.7. *Let M be a pseudoconvex, CR manifold of dimension $(2n+1)$ such that $\bar{\partial}_b$ has closed range in L^2 spaces for all degrees of forms. Let σ and $\hat{\sigma}$ be cutoff functions with disjoint supports and suppose that the σ - and $\hat{\sigma}$ -superlogarithmic properties on $(0, q_0)$ forms holds. Then*

$$\mathcal{G}_q \in C_{0,q,q,0}^\infty \left((V_0^\sigma \times V_0^{\hat{\sigma}}) \cup (V_0^{\hat{\sigma}} \times V_0^\sigma) \right) \quad \text{for all } q_0 \leq q \leq n - q_0.$$

If $q_0 = 1$ and $n \geq 2$, or $q_0 = n = 1$ with the extra assumption that G_0 is globally regular, then this conclusion also holds for $q = 0$ and $q = n$.

Proof. We choose $x_0 \in V_0^\sigma$ and $\hat{x}_0 \in V_0^{\hat{\sigma}}$ and two triplets of cutoff functions $\{\zeta_j\}_{j=0}^2$ and $\{\hat{\zeta}_j\}_{j=0}^2$ such that $\zeta_0 \prec \sigma \prec \zeta_1 \prec \zeta_2$, $\hat{\zeta}_0 \prec \hat{\sigma} \prec \hat{\zeta}_1 \prec \hat{\zeta}_2$, $\text{supp}(\zeta_2) \cap \text{supp}(\hat{\zeta}_2) = \emptyset$, and $x_0 \in V_0^{\zeta_0}$, $\hat{x}_0 \in V_0^{\hat{\zeta}_0}$. Since $\zeta_1 \varphi_{\hat{x}_0} = 0$, Theorem 4.1 implies that the desired inequality $\|\zeta_0 G_q \varphi_{\hat{x}_0}\|_{n+1+|\alpha|} < \infty$ holds if $\|\zeta_2 G_q^\delta \varphi_{\hat{x}_0}\|_{L^2} < \infty$ uniformly in $\delta > 0$. By the self-adjoint property of G^δ , the Hölder inequality and the estimate (4.24), we get

$$\begin{aligned} \|\zeta_2 G_q^\delta \varphi_{\hat{x}_0}\|_{L^2} &= \sup\{ |(\zeta_2 G_q^\delta \varphi_{\hat{x}_0}, v)_{L^2}| : \|v\|_{L^2} \leq 1 \} \\ &= \sup\{ |(\varphi_{\hat{x}_0}, \hat{\zeta}_0 G_q^\delta \zeta_2 v)_{L^2}| : \|v\|_{L^2} \leq 1 \} \\ &\leq \sup\{ \|\varphi_{\hat{x}_0}\|_{H^{-(n+1+|\hat{\alpha}|)}} \|\hat{\zeta}_0 G_q^\delta \zeta_2 v\|_{H^{n+1+|\hat{\alpha}|}} : \|v\|_{L^2} \leq 1 \} \\ &\leq \sup\{ \|\hat{\zeta}_0 G_q^\delta \zeta_2 v\|_{H^{n+1+|\hat{\alpha}|}} : \|v\|_{L^2} \leq 1 \}. \end{aligned} \quad (4.28)$$

We apply Theorem 3.10 for $u = G_q^\delta \zeta_2 v$ with system of cutoff functions $\{\hat{\zeta}_j\}$ to obtain

$$\begin{aligned} \|\hat{\zeta}_0 G_q^\delta \zeta_2 v\|_{H^{n+1+|\hat{\alpha}|}}^2 &\lesssim \|\hat{\zeta}_1 \zeta_2 v\|_{H^{n+1+|\hat{\alpha}|}}^2 + \|\hat{\zeta}_2 G_q^\delta \zeta_2 v\|_{L^2}^2 \\ &\leq \|G_q^\delta \zeta_2 v\|_{L^2}^2 \lesssim \|\zeta_2 v\|_{L^2}^2 \leq \|v\|^2 \leq 1. \end{aligned} \quad (4.29)$$

uniformly in δ . From (4.28) and (4.29), we get $\|\zeta_2 G_q^\delta \varphi_{\hat{x}_0}\|_{L^2}$ is uniformly bounded to δ . Therefore, $\|\zeta_0 G_q \varphi_{\hat{x}_0}\|_{n+1+|\alpha|} < \infty$ and hence by (4.27),

$$\mathcal{G}_q \in C_{0,q,q,0}^\infty(V_0^{\hat{\sigma}} \times V_0^\sigma).$$

We also notice that $\mathcal{G}_q$ is smoothness on symmetric sets of $M \times M$ since G_q is self adjoint. That means if $\mathcal{G}_q \in C_{0,q,q,0}^\infty(V_0^{\hat{\sigma}} \times V_0^\sigma)$ then $\mathcal{G}_q \in C_{0,q,q,0}^\infty(V_0^\sigma \times V_0^{\hat{\sigma}})$. This proves Theorem 4.7. $\square$

Theorem 4.8. *Let M be a pseudoconvex, CR manifold of dimension $(2n+1)$ such that $\bar{\partial}_b$ has closed range in L^2 spaces for all degrees of forms. Suppose that the σ -superlogarithmic property holds on $(0, q_0)$ -forms and G_q is globally regular for some $q_0 \leq q \leq n - q_0$. Then*

$$\mathcal{G}_q \in C_{0,q;q,0}^\infty \left(\left(V_0^\sigma \times (M \setminus \overline{\text{supp}(\sigma)}) \right) \cup \left((M \setminus \overline{\text{supp}(\sigma)}) \times V_0^\sigma \right) \right).$$

If $q_0 = q = 1$ (resp. $q_0 = 1, q = n$), this conclusion also holds for G_0 and (resp. G_n).

Proof. We choose $x_0 \in V_0^\sigma$ and $\hat{x}_0 \in M \setminus \overline{\text{supp}(\sigma)}$ and $\{\zeta_j\}_{j=0}^2$ such that $\zeta_0 \prec \sigma \prec \zeta_1 \prec \zeta_2$, and $\hat{x}_0 \in V_0^{\zeta_0}$, $\hat{x}_0 \notin \text{supp}(\zeta_1)$. This means $\zeta_1 \varphi_{\hat{x}_0} = 0$. By the global regularity of G_q , it follows for any multi-indices $\hat{\alpha}$ there exists $m_{\hat{\alpha}}$ such that

$$\|G_q v\|_{H^{n+1+|\hat{\alpha}|}}^2 \lesssim \|v\|_{H^{m_{\hat{\alpha}}}}^2 \quad \text{for all } v \in H_{0,q}^{m_{\hat{\alpha}}}(M); \quad (4.30)$$

and G_q can be extended to a continuous in the space of $H_{0,q}^{-\infty}(M)$. Thus we can estimate $\zeta_2 G_q \varphi_{\hat{x}_0}$ as follows

$$\begin{aligned} \|\zeta_2 G_q \varphi_{\hat{x}_0}\|_{H^{-m_{\hat{\alpha}}}} &= \sup\{ |(\zeta_2 G_q \varphi_{\hat{x}_0}, v)_{L^2}| : \|v\|_{H^{m_{\hat{\alpha}}}} \leq 1 \} \\ &= \sup\{ |(\varphi_{\hat{x}_0}, G_q \zeta_2 v)_{L^2}| : \|v\|_{L^2} \leq 1 \} \\ &\leq \sup\{ \|\varphi_{\hat{x}_0}\|_{H^{-(n+1+|\hat{\alpha}|)}} \|G_q \zeta_2 v\|_{H^{n+1+|\hat{\alpha}|}} : \|v\|_{H^{m_{\hat{\alpha}}}} \leq 1 \} \lesssim 1, \end{aligned}$$

where the last estimate follows by (4.30) and (4.24). Thus we can apply Theorem 3.8 for $u = G_q \zeta_2 \varphi_{\hat{x}_0}$, $s = n + 1 + |\hat{\alpha}|$, $s_o = m_{\hat{\alpha}}$ and $r = s_o + 2$, it follows

$$\|\zeta_0 G_q \varphi_{\hat{x}_0}\|_{H^{n+1+|\alpha|}}^2 \lesssim \|\zeta_1 \varphi_{\hat{x}_0}\|_{H^{n+1+|\alpha|}}^2 + \|\zeta_2 G_q \varphi_{\hat{x}_0}\|_{H^{-s_o}}^2 \lesssim 1.$$

By (4.27), it follows

$$\mathcal{G}_q \in C_{0,q;q,0}^\infty \left((M \setminus \overline{\text{supp}(\sigma)}) \times V_0^\sigma \right).$$

The proof of Theorem 4.8 follows by the symmetric of $\mathcal{G}_q$. $\square$

5. THE σ -SUPERLOGARITHMIC PROPERTY ON HYPERSURFACE MODELS

In this section we show that the σ -superlogarithmic property admits on the hypersurfaces model (1.10) and prove Theorem 1.9. Let M be the hypersurfaces passing the origin defined by

$$M = \left\{ (z, z_{n+1}) \in \mathbb{C}^n \times \mathbb{C} : \text{Im } z_{n+1} = \sum_k H_k \left(\sum_{l \in I_k} |h_l(z)|^2 \right) F_k \left(\sum_{l \in I_k} |\text{Re } h_l(z)|^2 \right) \right\}. \quad (5.1)$$

Here,

- (1) $H_k, F_k : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ are increasing and convex functions such that $H_k(0)F_k(0) = 0$ and. If $H_k(0) = 0$ (resp. $F_k(0) = 0$) it is added an extra assumption that $H_k(\delta)/\delta$ (resp. $F_k(\delta)/\delta$) is increasing. Here, we assume that $F_k(0) = 0$, otherwise the calculation is simpler.

- (2) h_l with $l = 1, \dots, N$ are holomorphic functions in $\mathbb{C}^n$ with an isolated zero at the origin and I_k 's are blocks of $(1, \dots, N)$ such that $\cup_k I_k \subset \{1, \dots, N\}$.

Theorem 5.1. *Let M be the hypersurface in $\mathbb{C}^{n+1}$ defined by (5.1). Then, if*

$$\lim_{\delta \rightarrow 0} \delta \ln (F_k(\delta^2)) = 0 \quad \text{for all } k, \quad (5.2)$$

then the origin of $\mathbb{C}^{n+1}$ is a weakly superlogarithmic point.

Proof. After the setting, the proof of this theorem is divided into three steps. In Step 1, we give a lower bounded of the Levi form of M by the $(1, 1)$ -forms

$$\mathcal{M}_l = \frac{H(|h_l|^2)F(|\operatorname{Re} h_l|^2)}{|\operatorname{Re} h_l|^2} \mathcal{L}_{|h_l|^2},$$

for all $l = 1, \dots, N$ where $H := \min_k H_k$ and $F := \min_k F_k$. In the language of the $(f, \mathcal{M})$ -property, the $(t^{1/2}, \mathcal{M}_l)$ -properties hold. Using Proposition 5.2, we interchange magnitude of f and $\mathcal{M}$ in the $(t^{1/2}, \mathcal{M}_l)$ -properties in Step 2. Finally, in Step 3 we construction a cutoff function σ in any small neighborhood in M of the origin x_o so that the σ -superlogarithmic property holds, and hence x_o is a weakly superlogarithmic point.

For setting, let

$$\rho := \sum_k \rho_k \quad \text{where} \quad \rho_k = H_k \left(\sum_{l \in I_k} |h_l|^2 \right) F_k \left(\sum_{l \in I_k} |\operatorname{Re} h_l|^2 \right).$$

We identify M with $\mathbb{C}^n \times \mathbb{R}$ so that the point $(z, \tau + \sqrt{-1}\rho) \in M$ corresponds to the point $(z, \tau) \in \mathbb{C}^n \times \mathbb{R}$. Set

$$L_j := \frac{\partial}{\partial z_j} + \sqrt{-1} \frac{\partial \rho}{\partial z_j}(z) \frac{\partial}{\partial \tau}, \quad T := \sqrt{-1} \frac{\partial}{\partial \tau}, \quad \theta := -\sqrt{-1} d\tau - \sum_{j=1}^n \left(\frac{\partial \rho}{\partial z_j} dz_j - \frac{\partial \rho}{\partial \bar{z}_j} d\bar{z}_j \right).$$

It is easy to verify that $L_j \in T^{1,0}M$ with its dual dz_j and θ is a contact form with its dual T .

Step 1. The Levi form of M is given by

$$d\theta = \sum_{i,j=1}^n \frac{\partial^2 \rho}{\partial z_i \partial \bar{z}_j} dz_i \wedge d\bar{z}_j = \frac{1}{2} (\partial_b \bar{\partial}_b - \bar{\partial}_b \partial_b) \rho = \mathcal{L}_\rho = \sum_k \mathcal{L}_{\rho_k}.$$

We use notation $|h^k|^2 = \sum_{j \in I_k} |h_j|^2$ and $|\operatorname{Re} h^k|^2 = \sum_{l \in I_k} |\operatorname{Re} h_l|^2$. We can check that

$$\mathcal{L}_{|h^k|^2} = \sum_{l \in I_k} \mathcal{L}_{|h_l|^2} = \sum_{l \in I_k} \partial_b h_l \wedge \bar{\partial}_b \bar{h}_l = 2\mathcal{L}_{|\operatorname{Re} h^k|^2}$$

and

$$\begin{aligned} \partial_b |h^k|^2 \wedge \bar{\partial}_b |\operatorname{Re} h^k|^2 + \partial_b |\operatorname{Re} h^k|^2 \wedge \bar{\partial}_b |h^k|^2 &= 2 \sum_{l, \bar{l} \in I_k} \operatorname{Re} h_l \partial_b h_l \wedge \operatorname{Re} h_{\bar{l}} \bar{\partial}_b \bar{h}_{\bar{l}} \\ &= 4 \partial_b |\operatorname{Re} h^k|^2 \wedge \bar{\partial}_b |\operatorname{Re} h^k|^2 \geq 0. \end{aligned}$$

Thus, by direct computation, we get

$$\begin{aligned} \mathcal{L}_{\rho_k} &= F_k \mathcal{L}_{H_k} + H_k \mathcal{L}_{F_k} + \partial_b H_k \wedge \bar{\partial}_b F_k + \partial_b F_k \wedge \bar{\partial}_b H_k \\ &= F_k \left(\dot{H}_k \mathcal{L}_{|h^k|^2} + \ddot{H}_k \partial_b |h^k|^2 \wedge \bar{\partial}_b |h^k|^2 \right) + H_k \left(\dot{F}_k \mathcal{L}_{|\operatorname{Re} h^k|^2} + \ddot{F}_k \partial_b |\operatorname{Re} h^k|^2 \wedge \bar{\partial}_b |\operatorname{Re} h^k|^2 \right) \\ &\quad + \dot{F}_k \dot{H}_k \left(\partial_b |h^k|^2 \wedge \bar{\partial}_b |\operatorname{Re} h^k|^2 + \partial_b |\operatorname{Re} h^k|^2 \wedge \bar{\partial}_b |h^k|^2 \right). \end{aligned} \quad (5.3)$$

From the last line of (5.3), it follows that $\mathcal{L}_{\rho_k} \geq \partial_b \rho_k \wedge \bar{\partial}_b \rho_k$; and from the second line of (5.3), it follows

$$\begin{aligned} \mathcal{L}_{\rho_k} &\geq (F_k \dot{H}_k + \frac{1}{2} \dot{F}_k H_k) \mathcal{L}_{|h^k|^2} \\ &\geq \frac{1}{2} \frac{H_k (|h^k|^2) F_k (|\operatorname{Re} h^k|^2)}{|\operatorname{Re} h^k|^2} \mathcal{L}_{|h^k|^2} \\ &\geq \frac{1}{2} \sum_{l \in I_k} \frac{H_k (|h_l|^2) F_k (|\operatorname{Re} h_l|^2)}{(|\operatorname{Re} h_l|^2)} \mathcal{L}_{|h_l|^2} \end{aligned} \quad (5.4)$$

where the inequalities follow by $\frac{F_k(\delta)}{\delta}$ is increasing. Consequently,

$$d\theta \gtrsim \sum_{l=1}^N \frac{H(|h_l|^2) F(|\operatorname{Re} h_l|^2)}{(|\operatorname{Re} h_l|^2)} \mathcal{L}_{|h_l|^2} + \partial_b \rho \wedge \bar{\partial}_b \rho. \quad (5.5)$$

where $H := \min_k H_k$ and $F := \min_k F_k$.

Step 2. In another words of the expression in (5.5), the $(f_1\text{-}\mathcal{M}_{l,1})$ -property holds for

$$f_1(t) = t^{1/2} \quad \text{and} \quad \mathcal{M}_{l,1} = \frac{H(|h_l|^2) F(|\operatorname{Re} h_l|^2)}{(|\operatorname{Re} h_l|^2)} \mathcal{L}_{|h_l|^2}.$$

Using Proposition 5.2 below, the $(f_2\text{-}\mathcal{M}_{l,2})$ -property holds for

$$f_2(t) = (F^*(t^{-1}))^{-1/2} \quad \text{and} \quad \mathcal{M}_{l,2} = H(|h_l|^2) \mathcal{L}_{|h_l|^2},$$

where F^* is denoted the inverse of F . By the hypothesis of F_k in (5.2) and $F(\delta) = \min_k F_k(\delta)$, we get $\lim_{\delta \rightarrow 0} \delta \ln F(\delta^2) = 0$. This implies $f_2 \gg \ln$. Denote $\tilde{H}(\delta) = \delta H(\delta)$.

Since $\mathcal{M}_{l,2} \geq \frac{\tilde{H}(|\operatorname{Re} h_l|^2)}{|\operatorname{Re} h_l|^2} \mathcal{L}_{|h_l|^2}$, we apply Proposition 5.2 again and obtain the $(f_3\text{-}\mathcal{M}_{l,3})$ -property holds for

$$f_3(t) = (\tilde{H}^*(F^*(t^{-1}))^{-1/2} \gg 1 \quad \text{and} \quad \mathcal{M}_{l,3} = \mathcal{L}_{|h_l|^2}.$$

Set

$$\mathcal{M}_2 := \sum_{l=1}^N H(|h_l|^2) \mathcal{L}_{|h_l|^2} \quad \text{and} \quad \mathcal{M}_3 := \sum_{l=1}^N \mathcal{L}_{|h_l|^2}.$$

Thus, the $(f_2\text{-}\mathcal{M}_2)$ - and $(f_3\text{-}\mathcal{M}_3)$ -properties holds. By Proposition 5.3 below, there exists $m \in \mathbb{N}$ such that $(f\text{-}Id)$ -property holds where $f(t) = (f_3(t))^{1/m}$ and $Id = \sum_{i=1}^n dz_i \wedge d\bar{z}_i$ the identity $(1, 1)$ -form of M . We conclude that in this step we have proven there exists a family of smooth real value function $\{\lambda_t\}$ such that $|T^m(\lambda_t)| \leq c_m e^{mt}$ such that

$$\mathcal{L}_{\lambda_t} + t d\theta \gtrsim f^2(t) Id + f_0^2(t) \ln^2 t \sum_{l=1}^N H(|h_l|^2) \mathcal{L}_{|h_l|^2} + t^{\frac{1}{2}} \partial_b \rho \wedge \bar{\partial}_b \rho. \quad (5.6)$$

where $f, f_0 \gg 1$ where $f_0 = f_2/\ln$.

Step 3. Let χ be a cutoff function in $\mathbb{R}$ such that $\chi(a) = \begin{cases} 1 & \text{if } |a| \leq 1, \\ 0 & \text{if } |a| \geq 2. \end{cases}$. Denote by $\chi_\epsilon(a) = \chi(a/\epsilon)$. We set

$$h_0 := \tau \quad \text{and} \quad \sigma_\epsilon := \prod_{l=0}^N \chi_\epsilon(|h_l|^2).$$

Since the common zero of h_l , for $l = 1, \dots, N$, is at only the origin of $\mathbb{C}^n$. Thus σ_ϵ is a cutoff function on M . Furthermore, for any small neighborhood U in M of x_o (the origin of M) there exists $\epsilon > 0$ such that $\text{supp}(\sigma_\epsilon) \subset U$ and $\sigma_\epsilon = 1$ in another neighborhood of x_o . In order to prove that x_o is the weakly superlogarithmic point, we shall show that the σ_ϵ -superlogarithmic property holds in U . Before giving the upper bounds of $\partial_b \sigma_\epsilon \wedge \bar{\partial}_b \sigma_\epsilon$ and $\mathcal{L}_{\sigma_\epsilon}$, we compute

$$\mathcal{L}_{|h_0|^2} = 2\partial_b h_0 \wedge \bar{\partial}_b h_0 + 2h_0 \mathcal{L}_{h^0} = 2\partial_b \tau \wedge \bar{\partial}_b \tau + 2\tau \mathcal{L}_\tau.$$

We have

$$\partial_b \tau = \sum_{j=1}^n L_j \tau dz_j = \sqrt{-1} \sum_{j=1}^n \frac{\partial \rho}{\partial z_j} dz_j = \sqrt{-1} \partial_b \rho.$$

Thus, $\bar{\partial}_b \tau = -\sqrt{-1} \bar{\partial}_b \rho$ and hence

$$\partial_b \tau \wedge \bar{\partial}_b \tau = \partial_b \rho \wedge \bar{\partial}_b \rho \underset{\text{by (5.5)}}{\lesssim} \mathcal{L}_\rho = d\theta.$$

$$\mathcal{L}_\tau = \frac{1}{2} (\partial_b \bar{\partial}_b \tau - \bar{\partial}_b \partial_b \tau) = \frac{1}{2} \sum_{ij=1}^n \left(-\sqrt{-1} \frac{\partial^2 \rho}{\partial z_i \partial \bar{z}_j} dz_i \wedge d\bar{z}_j - \sqrt{-1} \frac{\partial^2 \rho}{\partial \bar{z}_i \partial z_j} d\bar{z}_i \wedge dz_j \right) \equiv 0.$$

Thus $\mathcal{L}_{|h_0|^2} \lesssim d\theta$. We are ready to estimate $\partial_b \sigma_\epsilon \wedge \bar{\partial}_b \sigma_\epsilon$ and $\mathcal{L}_{\sigma_\epsilon}$

$$\partial_b \sigma_\epsilon = \sum_{l=0}^N \left(\prod_{\hat{l} \neq l} \chi(|h_{\hat{l}}|^2) \right) \left(\dot{\chi}(|h_l|^2) \partial_b |h_l|^2 \right)$$

and

$$\begin{aligned} \mathcal{L}_{\sigma_\epsilon} = & \sum_{l=0}^N \left\{ \left(\prod_{\hat{l} \neq l} \chi(|h_{\hat{l}}|^2) \right) \left(\dot{\chi}(|h_l|^2) \mathcal{L}_{|h_l|^2} + \ddot{\chi}(|h_l|^2) \partial |h_l|^2 \wedge \bar{\partial}_b |h_l|^2 \right) \right\} \\ & + \sum_{k,l=0, k \neq l}^N \left\{ \left(\prod_{\hat{l} \neq k,l} \chi(|h_{\hat{l}}|^2) \right) \left(\dot{\chi}(|h_k|^2) \dot{\chi}(|h_l|^2) \partial |h_k|^2 \wedge \bar{\partial}_b |h_l|^2 \right) \right\}. \end{aligned}$$

Thus, it follows

$$\begin{aligned} \ln t |\langle \mathcal{L}_{\sigma_\epsilon} \lrcorner u, u \rangle| + \ln^2 t |\partial_b \sigma_\epsilon \lrcorner u|^2 & \lesssim \ln^2 t \sum_{l=0}^N I_{\text{supp}(\dot{\chi}_\epsilon(|h_l|^2))} |\langle \mathcal{L}_{|h_l|^2} \lrcorner u, u \rangle| \\ & \leq \ln^2 t \left(\langle d\theta \lrcorner u, u \rangle + (H(\epsilon^2))^{-1} \ln^2 t \sum_{l=1}^N H(|h_l|^2) \langle \mathcal{L}_{|h_l|^2} \lrcorner u, u \rangle \right) \end{aligned} \quad (5.7)$$

for any $t \geq 1$ and $u \in C_{0,1}^\infty(M)$. where the second inequality follows by the fact that

$$I_{\text{supp}(\dot{\chi}_\epsilon(|h_l|^2))} \leq I_{\{x \in M : \epsilon \leq |h_l(x)| \leq 2\epsilon\}} \leq H(|\epsilon|^2)^{-1} H(|h_l|^2).$$

Combining (5.6) and (5.7), we can conclude that the σ_ϵ -superlogarithmic property holds for any $\epsilon > 0$. $\square$

The proof of Theorem 5.1 is complete but we have skipped a crucial technical point in Step 2 of the proof that we face now. First, the interchange sizes of the rate f and the multiplier $\mathcal{M}$ is expressed by the following proposition.

Proposition 5.2. *Let M be the hypersurface defined by (5.1) and h be a holomorphic function in $\mathbb{C}^n$. Assume that the $(f_1, \mathcal{M}_1)$ -property holds with a rate $1 \ll f_1 \lesssim t^{1/2}$ and multiplier*

$$\mathcal{M}_1 \geq \frac{H(|h|^2) F(|\text{Re } h|^2)}{|\text{Re } h|^2} \mathcal{L}_{|h|^2}.$$

Then the $(f_2, \mathcal{M}_2)$ -property holds for $f_2(t) = (F^*(f_1(t)^{-2}))^{-1/2}$ and $\mathcal{M}_2 = H(|h|^2) \mathcal{L}_{|h|^2}$. Here $F, H : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ are smooth increasing, convex functions with $F(0) = 0$, $\frac{F(\delta)}{\delta} \searrow 0$; and F^* is denoted the inverse of F .

Proof. Assume that for any $t \geq 1$ there exists a function $\lambda_{t,1}$ such that

$$|T^s(\lambda_{t,1})| \leq c_{s,1} e^{ts/2}, \quad \text{for all } s \in \mathbb{N} \text{ and } \mathcal{L}_{\lambda_{t,1}} + td\gamma \geq f_1(t)^2 \mathcal{M}_1.$$

Now we construct $\lambda_{t,2}$ such that

$$|T^s(\lambda_{t,2})| \leq c_{s,2}e^{ts/2}, \quad \text{for all } s \in \mathbb{N} \text{ and } \mathcal{L}_{\lambda_{t,2}} + td\gamma \geq f_2(t)^2\mathcal{M}_2. \quad (5.8)$$

We define

$$\tilde{\lambda}_t(z, \tau) = \lambda_t(|\operatorname{Re} h(z)|^2)H(|h(z)|^2) \quad \text{where} \quad \lambda_t(a) := \left(1 - e^{-\frac{1}{2}f_2^2(t)a}\right).$$

Since $\dot{\lambda}_t(a) = \frac{1}{2}f_2^2(t)\lambda_t(a) \geq 0$ and $\ddot{\lambda}_t(a) = -\frac{1}{4}f_2^4(t)\lambda_t(a)$, it follows

$$\begin{aligned} \mathcal{L}_{\tilde{\lambda}_t} &= H\mathcal{L}_{\lambda_t} + \lambda_t\mathcal{L}_H + \partial_b(H) \wedge \bar{\partial}_b(\lambda_t) + \partial_b(\lambda_t) \wedge \bar{\partial}_b(H) \\ &\geq H\left(\dot{\lambda}_t + |\operatorname{Re} h|^2\dot{\lambda}_t\right)\partial_b h \wedge \bar{\partial}_b \bar{h} \\ &\geq \frac{1}{4}f_2^2(t)He^{-\frac{1}{2}f_2^2(t)|\operatorname{Re} h|^2}\left\{\left(2 - f_2^2(t)|\operatorname{Re} h|^2\right)\partial_b h \wedge \bar{\partial}_b \bar{h}\right\} \end{aligned} \quad (5.9)$$

If $f_2|\operatorname{Re} h| \leq 1$, then $\exp\left(-\frac{1}{2}f_2^2(t)h^2\right) \geq e^{-1/2}$ uniformly in t and hence

$$\mathcal{L}_{\tilde{\lambda}_t} \geq \frac{1}{4e^{1/2}}f_2^2(t)H(|h|^2)\partial_b h \wedge \bar{\partial}_b \bar{h} \geq \frac{1}{8e^{1/2}}f_2^2(t)\mathcal{M}_2.$$

Otherwise, assume that $f_2|\operatorname{Re} h| \geq 1$. Using our assumption that $F(\delta)/\delta$ increasing for $|\operatorname{Re} h| \geq f_2^{-1}$ and the definition of $f_2(t) = (F^*(f_1(t))^{-2})^{-1/2}$, we get

$$f_1^2(t)\frac{F(|\operatorname{Re} h|^2)}{|\operatorname{Re} h|^2} \geq f_1^2(t)\frac{F((f_2(t))^{-2})}{(f_2(t))^{-2}} = f_2^2(t).$$

In this case, $\mathcal{L}_{\tilde{\lambda}}$ can get negative; however, by using the fact that $\min_{a \geq 1} \left\{(2-a)e^{-\frac{1}{2}a}\right\} = -2e^{-2}$ at $a = 4$ for $a = f_2(t)|\operatorname{Re} h| \geq 1$, we have

$$\mathcal{L}_{\tilde{\lambda}_t} \geq -\frac{1}{2e^2}f_2^2(t)\mathcal{M}_2$$

Thus

$$f_1^2(t)\mathcal{M}_1 + \mathcal{L}_{\tilde{\lambda}_t} \geq \left(1 - \frac{1}{2e^2}\right)f_2(t)^2\mathcal{M}_2.$$

In both cases, we have proved

$$f_1^2(t)\mathcal{M}_1 + \mathcal{L}_{\tilde{\lambda}_t} \geq \frac{1}{8e^{1/2}}f_2(t)^2\mathcal{M}_2.$$

Finally, we set $\lambda_{t,2} = \lambda_{t,1} + \tilde{\lambda}_t$. It is easy to check that both $|T^s(\lambda_{t,2})| \leq c_{s,2}e^{ts/2}$ for some $c_{s,2} > 0$. Therefore, $\lambda_{t,2}$ satisfies 5.8 and hence Proposition 5.2 is proven. $\square$

Proposition 5.3. *Let M be the hypersurface defined by (5.1) and $h_1, \dots, h_N$ be holomorphic functions defined in a neighborhood of the origin in $\mathbb{C}^n$ such that and*

- (1) *the common zero of h_l , $l = 1, \dots, N$ is isolated at the origin of $\mathbb{C}^n$;*
- (2) *the $(f\text{-}\mathcal{L}_{|h_l^2|})$ -properties hold for all $l = 1, \dots, N$ with some $f \gg 1$.*

Then, there exist an integer m such that the property $(\sqrt[m]{f}\text{-Id})$ holds.

Following the guidelines of Kohn in [Koh79, Koh00], we consider the ideas of germ of holomorphic function $\mathcal{J}_p$ defined inductively as follows. If $\mathcal{G}$ is a P -tuple $(g_1, \dots, g_P)$ is of germs of holomorphic functions in $\mathbb{C}^n$ let $\mathcal{B}(\mathcal{G})$ denote the $n \times (N + P)$ matrix

$$\mathcal{B}(\mathcal{G}) = \begin{pmatrix} \frac{\partial h_1}{\partial z_1} & \frac{\partial h_1}{\partial z_2} & \cdots & \frac{\partial h_1}{\partial z_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial h_N}{\partial z_1} & \frac{\partial h_N}{\partial z_2} & \cdots & \frac{\partial h_N}{\partial z_n} \\ \frac{\partial g_1}{\partial z_1} & \frac{\partial g_1}{\partial z_2} & \cdots & \frac{\partial g_1}{\partial z_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial g_P}{\partial z_1} & \frac{\partial g_P}{\partial z_2} & \cdots & \frac{\partial g_P}{\partial z_n} \end{pmatrix}.$$

Denote by $\det \mathcal{B}(\mathcal{G})$ the ideal generated by all determinants of all the $n \times n$ submatrices of $\mathcal{B}(\mathcal{G})$. Then we define

$$\mathcal{J}_0 = \sqrt{\det \mathcal{B}(\{0\})}$$

and for $p \geq 1$

$$\mathcal{J}_p = \sqrt{(\mathcal{J}_{p-1}, \det \mathcal{B}(\mathcal{J}_{p-1}))}.$$

Here, $\sqrt{\mathcal{J}}$ is a ideal of holomorphic functions such that if $g \in \sqrt{\mathcal{J}}$ then there exists $\tilde{g} \in \mathcal{J}$ and $m \in \mathcal{N}$ such that $|g|^m \leq |\tilde{g}|$, where $\mathcal{J}$ is a ideal of germs of holomorphic functions.

For the proof of Proposition 5.3, we also need the following lemma.

Lemma 5.4. *Let g be a holomorphic function and $f \gg 1$. Then*

$$\text{The } (f|g|^{2m} \text{Id})\text{-property} \implies \text{the } (f^{1/m}|g|^2 \text{Id})\text{-property}.$$

Proof. The proof follows immediately by the fact that $\mathcal{L}_{|z|^2} = \text{Id}$ (i.e., the (1-Id) -property holds) and

$$f^{2/m}(t)|g(z)|^2 \leq [f^{2/m}(t)|g(z)|^2]^m + 1 = f^2(t)|g|^{2m} + 1.$$

□

We are ready to prove Proposition 5.3.

Proof. It was proved in [Koh79, Theorem 7.13] that $1 \in \mathcal{J}_p$ for some integer p if and only if the common zero of the functions $h_1, \dots, h_N$ is the origin of $\mathbb{C}^n$. Thus, we only need to prove that if $g \in \mathcal{J}_p$ then there exists an integer m_p such that $(f^{1/m_p}|g|^2 \text{Id})$ -property holds. We prove this statement by the inductive method. The argument for checking the first step $p = 0$, by using the hypothesis that the $(f\mathcal{L}_{|h_j|^2})$ -property holds for all $j = 1, \dots, N$, is similar for a general step p so we omit here the step $p = 0$. Now we assume inductively that the $(f^{\frac{1}{m_{p-1}}}|g|^2 \text{Id})$ holds for all $\hat{g} \in \mathcal{J}_{p-1}$. By Proposition 5.2, the $(f^{\frac{1}{2m_{p-1}}}\mathcal{L}_{|\hat{g}|^2})$ -property holds for all $\hat{g} \in \mathcal{J}_{p-1}$ and hence by the definition of $\det \mathcal{B}(\mathcal{J}_{p-1})$, we get $(f^{\frac{1}{2m_{p-1}}}|g|^2 \text{Id})$ -property holds for all $\tilde{g} \in (\mathcal{J}_{p-1}, \det \mathcal{B}(\mathcal{J}_{p-1}))$. Now taking $g \in \mathcal{J}_p$

there exist an integer $\tilde{m}_q$ and $\tilde{g} \in (\mathcal{J}_{p-1}, \det \mathcal{B}(\mathcal{J}_{p-1}))$ such that $|g|^{\tilde{m}_p} \leq |\tilde{g}|$. By Lemma 5.4, the $(f^{\frac{1}{m_p}} - |g|^2 \text{Id})$ -property holds for $m_p := 2m_{p-1}\tilde{m}_p$. This proves Proposition 5.3. $\square$

Now we are ready for the proof of Theorem 1.9.

Theorem 1.9. By the argument in Theorem 5.1, we can prove that any $x \in M$ is a weakly superlogarithmic point. In order to use Theorems 1.3 and 1.5, we have to show that $\bar{\partial}_b$ has closed range in L^2 for all degree of forms and G_0 is globally regular in the case $n = 1$. Up to the author's knowledge, the closed range of $\bar{\partial}_b$ and global regularity of G_0 have been established only on compact CR manifolds (cf. [Bar12, Koh86, KN06, Nic06, Sha85, SZ15]) nevertheless M is not a compact CR manifold. Since M is a hypersurface in $\mathbb{C}^{n+1}$, we can compactify M by [McN92, Proposition 2.1] as follows. Let V be an open set in M , there exists a smooth, compact, pseudoconvex hypersurface $\tilde{M}$ in $\mathbb{C}^{n+1}$ satisfying the following properties:

- $V \subset M \cap \tilde{M}$,
- all points in $\tilde{M} \setminus M$ are points of strong pseudoconvexity,
- the relative boundary S of $\tilde{M} \cap M$ and $\tilde{M} \setminus M$ is the intersection of M with a sphere centered at the origin of $\mathbb{C}^{n+1}$.

Thus, the $\bar{\partial}_{b,\tilde{M}}$ has closed range where the subscript $\tilde{M}$ means the operator of $\tilde{M}$. Analogously to [KZ12, Proposition 4.1], we have $G_{q,\tilde{M}}$ is globally exactly regular for all $0 \leq q \leq n$. Indeed, since the compactness estimate for the system $(\bar{\partial}_{b,M}, \bar{\partial}_{b,M}^*)$ holds on any open set in M (see Step 2 in the proof of Theorem 5.1), similar to the argument in the proof of [KZ12, Proposition 4.1], we can obtain the (global) compactness estimate for the system $(\bar{\partial}_{b,\tilde{M}}, \bar{\partial}_{b,\tilde{M}}^*)$ and hence the global regularity of $G_{q,\tilde{M}}$ follows (see [Rai10, KPZ12, Str12, SZ15]). Now we can apply Theorems 1.3 and 1.5, we obtain $\square_{b,\tilde{M}}$ is locally hypoelliptic. On local set $V \subset M \cap \tilde{M}$, the Kohn-Laplacians $\square_{b,M}$ and $\square_{b,\tilde{M}}$ coincide. Therefore, $\square_{b,M}$ is local hypoelliptic on V . This completes the proof of Theorem 1.9. $\square$

APPENDIX

In this appendix, we give estimates of commutators to support the work in §2. The following lemma has been used in the proof of Proposition 2.4

Lemma 5.5. *Let $\lambda := \lambda_{k,s}$ be a real-valued C^∞ function defined on an open set U of M such that $|\lambda| \leq c_{0,s}k$ and $|T^m(\lambda)| \leq c_{m,s}e^{\frac{mk}{2}}$ for any $m \in \mathbb{N}$. Then, for any $p, r \in \mathbb{N}$ and $\alpha \in \mathbb{R}$, there exists a constant $c_{s,r,p,\alpha}$ independent of k such that*

$$e^{\alpha k} \|T^p[e^{-\frac{\lambda}{2}} \zeta, \tilde{\Gamma}_k^+] \Gamma_k \Psi^+ \zeta u\|_{L^2} \leq c_{s,r,p,\alpha} \|\zeta u\|_{H^{-r}}$$

holds for all $u \in H^{-s_o}(M)$ with $r \geq s_o$. Here, $\mathcal{S}(\tilde{\Gamma}_k^+) \succ \mathcal{S}(\Gamma_k \Psi^+)$.

We first remark that the supports of the symbols of $[e^{-\frac{\lambda}{2}}\zeta, \tilde{\Gamma}_k^+]$ and $\Gamma_k\Psi^+$ are disjoint, the operator $[e^{-\frac{\lambda}{2}}\zeta, \tilde{\Gamma}_k^+]\Gamma_k\Psi^+$ is order $-\infty$. This implies $\|T^p[e^{-\frac{\lambda}{2}}\zeta, \tilde{\Gamma}_k^+]\Gamma_k\Psi^+\zeta u\|^2$ can be controlled by any negative Sobolev norm with multiplying by a constant. However this constant might depend on k . The Lemma 5.5 is to verify that the upper bounded of $e^{\alpha k}\|T^p[e^{-\frac{\lambda}{2}}\zeta, \tilde{\Gamma}_k^+]\Gamma_k\Psi^+\zeta u\|^2$ is independent on k .

Proof. Let

$$K(\xi_{2n+1}, \eta_{2n+1}) = \xi_{2n+1}^p [\tilde{\gamma}(e^{-k}\xi_{2n+1}) - \tilde{\gamma}(e^{-k}\eta_{2n+1})] \mathcal{F}_{2n+1}\{e^{-\frac{\lambda}{2}}\zeta\}(\xi_{2n+1} - \eta_{2n+1})\eta_{2n+1}^r$$

on support of $\gamma(e^{-k}\eta_{2n+1})\mathcal{F}_{2n+1}(\Psi^+\zeta v)(\eta_{2n+1})$. From Taylor's theorem, we have for any $m \geq 1$ there exists $a_m \in [0, 1]$ such that

$$\begin{aligned} \tilde{\gamma}(e^{-k}\xi_{2n+1}) - \tilde{\gamma}(e^{-k}\eta_{2n+1}) &= \sum_{j=1}^{m-1} \frac{1}{j!} e^{-jk} (\xi_{2n+1} - \eta_{2n+1})^j \tilde{\gamma}^{(j)}(e^{-k}\eta_{2n+1}) \\ &\quad + \frac{1}{m!} e^{-mk} (\xi_{2n+1} - \eta_{2n+1})^m \tilde{\gamma}^{(m)}(a_m e^{-k}\xi_{2n+1} + (1 - a_m)e^{-k}\eta_{2n+1}). \end{aligned} \quad (5.10)$$

On support of $\gamma(e^{-k}\eta_{2n+1})$, the summation $\sum_{j=1}^{m-1} \bullet$ is zero since $\mathcal{S}(\tilde{\Gamma}_k^+) \succ \mathcal{S}(\Gamma_k\Psi^+)$ and hence

$$\begin{aligned} |K(\xi_{2n+1}, \eta_{2n+1})| &\leq c_m e^{-mk} |\eta_{2n+1}^r| \left| \xi_{2n+1}^p (\xi_{2n+1} - \eta_{2n+1})^m \mathcal{F}_{2n+1}\{e^{-\frac{\lambda}{2}}\zeta\}(\xi_{2n+1} - \eta_{2n+1}) \right| \\ &\leq c_{m,p} e^{-mk} |\eta_{2n+1}^r| \left[\underbrace{(\xi_{2n+1} - \eta_{2n+1})^{m+p} \mathcal{F}_{2n+1}\{e^{-\frac{\lambda}{2}}\zeta\}(\xi_{2n+1} - \eta_{2n+1})}_{I_{m+p}} \right] \\ &\quad + \underbrace{|\eta_{2n+1}^{p+r} (\xi_{2n+1} - \eta_{2n+1})^m \mathcal{F}_{2n+1}\{e^{-\frac{\lambda}{2}}\zeta\}(\xi_{2n+1} - \eta_{2n+1})|}_{I_m}. \end{aligned} \quad (5.11)$$

For $l = m + p$ or m , we rewrite I_l as

$$\begin{aligned} I_l &= (|\xi_{2n+1} - \eta_{2n+1}|^2 + 1)^{-1} \mathcal{F}_{2n+1}\{[(\xi_{2n+1} - \eta_{2n+1})^{l+2} + (\xi_{2n+1} - \eta_{2n+1})^l]e^{-\frac{\lambda}{2}}\zeta\}(\xi_{2n+1} - \eta_{2n+1}) \\ &= (|\xi_{2n+1} - \eta_{2n+1}|^2 + 1)^{-1} \mathcal{F}_{2n+1}\{(e^{-\frac{\lambda}{2}}\zeta)^{(l+2)} + (e^{-\frac{\lambda}{2}}\zeta)^{(l)}\}(\xi_{2n+1} - \eta_{2n+1}) \\ &= (|\xi_{2n+1} - \eta_{2n+1}|^2 + 1)^{-1} \left[\mathcal{F}_{2n+1}\{(e^{-\frac{\lambda}{2}}\zeta)^{(l+2)}\}(\xi_{2n+1} - \eta_{2n+1}) + \mathcal{F}_{2n+1}\{(e^{-\frac{\lambda}{2}}\zeta)^{(l)}\}(\xi_{2n+1} - \eta_{2n+1}) \right]. \end{aligned} \quad (5.12)$$

On other hand,

$$\begin{aligned} |\mathcal{F}_{2n+1}\{(e^{-\frac{\lambda}{2}}\zeta)^l\}(\xi_{2n+1} - \eta_{2n+1})| &= \left| \int_{\mathbb{R}} e^{-i(\xi_{2n+1} - \eta_{2n+1})x_{2n+1}} T^l(e^{-\frac{\lambda}{2}}\zeta) dx_{2n+1} \right| \\ &\leq \int_{\mathbb{R}} |T^l(e^{-\frac{\lambda}{2}}\zeta)| dx_{2n+1} \\ &\leq c_{l,s} e^{\frac{lk}{2} + c_{0,s}k}, \end{aligned} \quad (5.13)$$

where the last inequality follows by the assumption on $\lambda := \lambda_{k,s}$. Thus, $|I_l| \leq c_{l,s} e^{\frac{k(l+2)}{2} + c_{0,s}k}$. Note that on support of $\gamma(e^{-k}\eta_{2n+1})$, $|\eta_{2n+1}| \leq e^k$. Therefore,

$$|K(\xi_{2n+1}, \eta_{2n+1})| \leq c_{m,p,s} (|\xi_{2n+1} - \eta_{2n+1}|^2 + 1)^{-1} e^{-mk + \frac{(m+2)k}{2} + (r+p+c_{0,s})k},$$

holds for any $m \in \mathbb{N}$. Fix $\alpha \in \mathbb{R}$ choose m sufficiently large, we got

$$|K(\xi_{2n+1}, \eta_{2n+1})| \leq c_{\alpha,p,r,s} e^{-\alpha k} (|\xi_{2n+1} - \eta_{2n+1}|^2 + 1)^{-1}.$$

This boundedness of the kernel $K(\xi_{2n+1}, \eta_{2n+1})$ implies that

$$e^{\alpha k} \|T^p [e^{-\frac{\lambda k}{2}} \zeta, \tilde{\Gamma}_k^+] \Gamma_k \Psi^+ \zeta u\|_{L^2} \leq c_{\alpha,p,r,s} \|T^{-r} \Gamma_k \Psi^+ \zeta u\|_{L^2} \leq c_{\alpha,p,r,s} \|\zeta u\|_{H^{-r}}.$$

This completes the proof of this lemma. $\square$

The following lemma has been used to estimate the commutators

$$\sum_k \|e^{ks\sigma} [\bar{\partial}_b, \zeta \Gamma_k \Psi^+ \zeta] u\|_{L^2}^2 + \|e^{ks\sigma} [\bar{\partial}_b^*, \zeta \Gamma_k \Psi^+ \zeta] u\|_{L^2}^2$$

in Lemma 2.5 in §2.

Lemma 5.6. *Let X be a differential operator of order one with smooth coefficients and $u \in H^{-s_0}(M)$. Then, the following holds uniformly in k .*

$$(i) \quad \|e^{ks\sigma} [X, \zeta \Gamma_k \Psi^\pm \zeta] u\|_{L^2}^2 \leq c_s \left(\|e^{(k-1)s\sigma} \zeta \Gamma_{k-1} \Psi^\pm \zeta u\|_{L^2}^2 + \|e^{(k+1)s\sigma} \zeta \Gamma_{k+1} \Psi^\pm \zeta u\|_{L^2}^2 \right) \\ + c \left(\|\tilde{\Gamma}_k \tilde{\zeta} u\|_{L^2}^2 + e^{2ks} \|\tilde{\Gamma}_k \tilde{\Psi}^0 \zeta u\|_{L^2}^2 \right);$$

$$(ii) \quad \|X e^{ks\sigma} \zeta \Gamma_k \Psi^\pm \zeta u\|_{L^2}^2 \leq c \left(\|e^{ks\sigma+k} \zeta \Gamma_k \Psi^\pm \zeta u\|_{L^2}^2 + e^{2k} \|\Gamma_k \zeta u\|_{L^2}^2 \right).$$

Here we recall that ζ is a cutoff function such that $\zeta \prec \tilde{\zeta}$; $\tilde{\Psi}^0$ is defined in §2.2 such that $\mathcal{S}([\Psi^\pm, \alpha]) \prec \mathcal{S}(\tilde{\Psi}^0)$; and $\tilde{\Gamma}_k$ is a pseudodifferential operator such that $\mathcal{S}(\Gamma_k) \prec \mathcal{S}(\tilde{\Gamma}_k)$.

Proof. For the proof of (i), we first use the Jacobi identity to get

$$\|e^{ks\sigma} [X, \zeta \Gamma_k \Psi^\pm \zeta] u\|_{L^2}^2 \leq 4 \left(\|e^{ks\sigma} \zeta [X, \Gamma_k] \Psi^\pm \zeta u\|_{L^2}^2 + \|e^{ks\sigma} \zeta \Gamma_k [X, \Psi^\pm] \zeta u\|_{L^2}^2 \right. \\ \left. + \|e^{ks\sigma} [X, \zeta] \Gamma_k \Psi^\pm \zeta u\|_{L^2}^2 + \|e^{ks\sigma} \zeta \Gamma_k \Psi^\pm [X, \zeta] u\|_{L^2}^2 \right). \quad (5.14)$$

Since $\text{supp}(\dot{\zeta}) \cap \text{supp}(\sigma) = \emptyset$, the two terms in the second line of (5.14) are bounded by $\|\tilde{\Gamma}_k \tilde{\zeta} u\|_{L^2}^2$. By the choice of $\tilde{\Psi}^0$ and $\tilde{\Gamma}_k$, it follows $\mathcal{S}(\Gamma_k [X, \Psi^\pm]) \prec \mathcal{S}(\tilde{\Gamma}_k \tilde{\Psi}^0)$, thus the last term in the first line of (5.14) can be bounded as follows

$$\|e^{ks\sigma} \zeta \Gamma_k [X, \Psi^\pm] \zeta u\|_{L^2}^2 \leq e^{2ks} \|\Gamma_k [X, \Psi^\pm] \tilde{\Psi}^0 \zeta u\|_{L^2}^2 \leq c e^{2ks} \|\tilde{\Gamma}_k \tilde{\Psi}^0 \zeta u\|_{L^2}^2. \quad (5.15)$$

The first term in the RHS. of the first line of (5.14) is treated as follows. Since $(\gamma_{k+1} + \gamma_{k-1}) = 1$ on the support of $\dot{\gamma}_k$, we have

$$\begin{aligned} \|e^{ks\sigma}\zeta[X, \Gamma_k]\Psi^\pm\zeta u\|_{L^2}^2 &= \|e^{ks\sigma}\zeta[X, \Gamma_k]\tilde{\Psi}^\pm(\Gamma_{k-1} + \Gamma_{k+1})\Psi^\pm\zeta u\|_{L^2}^2 \\ &\leq \| [X, \Gamma_k]\tilde{\Psi}^\pm e^{ks\sigma}\zeta(\Gamma_{k-1} + \Gamma_{k+1})\Psi^\pm\zeta u\|_{L^2}^2 \\ &\quad + \|e^{ks\sigma}, [X, \Gamma_k]\tilde{\Psi}^\pm\zeta(\Gamma_{k-1} + \Gamma_{k+1})\Psi^\pm\zeta u\|_{L^2}^2 \\ &\quad + \|e^{ks\sigma}[\zeta, [X, \Gamma_k]\tilde{\Psi}^\pm](\Gamma_{k-1} + \Gamma_{k+1})\Psi^\pm\zeta u\|_{L^2}^2. \end{aligned} \quad (5.16)$$

The term in the last line of (5.16) is bounded by $\|\tilde{\Gamma}_k\tilde{\zeta}u\|_{L^2}^2$ since $\text{supp}(\dot{\zeta}) \cap \text{supp}(\sigma) = \emptyset$. Since the symbol of $[X, \Gamma_k]\tilde{\Psi}^\pm$ is uniformly bounded by a constant, the term in second line of (5.16) is bounded by $\|e^{ks\sigma}\zeta(\Gamma_{k-1} + \Gamma_{k+1})\Psi^\pm\zeta u\|_{L^2}^2$. We also notice that the support of $\mathcal{S}([X, \Gamma_k]\tilde{\Psi}^\pm)$ is a subset of $\text{supp}(\dot{\gamma}_k\tilde{\psi}^+)$ and hence

$$\| [e^{ks\sigma}, [X, \Gamma_k]\tilde{\Psi}^\pm]v\|_{L^2}^2 \leq c_s k^2 e^{-2k} \|e^{ks\sigma}v\|_{L^2}^2 \leq c_s \|e^{ks\sigma}v\|_{L^2}^2.$$

Thus, the term in the third line of (5.16) is also bounded by $\|e^{ks\sigma}\zeta(\Gamma_{k-1} + \Gamma_{k+1})\Psi^\pm\zeta u\|_{L^2}^2$. This proves the first part of Lemma 5.6.

For the proof of (ii), we have

$$\begin{aligned} \|Xe^{ks\sigma}\zeta\Gamma_k\Psi^\pm\zeta u\|_{L^2}^2 &\lesssim \|\Lambda\tilde{\zeta}e^{ks\sigma}\zeta\tilde{\Gamma}_k\tilde{\Psi}^\pm\Gamma_k\Psi^\pm\zeta u\|_{L^2}^2 \\ &\lesssim \|\Lambda\tilde{\Gamma}_k\tilde{\Psi}^\pm\tilde{\zeta}e^{ks\sigma}\zeta\Gamma_k\Psi^\pm\zeta u\|_{L^2}^2 + \|\Lambda[\tilde{\zeta}e^{ks\sigma}\zeta, \tilde{\Gamma}_k\tilde{\Psi}^\pm]\Gamma_k\Psi^\pm\zeta u\|_{L^2}^2. \end{aligned} \quad (5.17)$$

The symbol $\mathcal{S}(\Lambda\tilde{\Gamma}_k\tilde{\Psi}^\pm)$ is bounded by e^k since, so the first term in the second line of (5.17) is bounded by $\|e^{ks\sigma+1}\zeta\Gamma_k\Psi^\pm\zeta u\|_{L^2}^2$. Similar to the argument in the previous part, the second term in the second line of (5.17) is bounded by $k^2\|e^{ks\sigma}\zeta\Gamma_k\Psi^\pm\zeta u\|_{L^2}^2 + e^{2k}\|\Gamma_k\zeta u\|_{L^2}^2$. This proves (ii) and then the proof of Lemma 5.6 is complete. $\square$

REFERENCES

- [Bar12] L. Baracco. The range of the tangential Cauchy-Riemann system to a CR embedded manifold. *Invent. Math.*, 190(2):505–510, 2012.
- [BKZ12] L. Baracco, T.V. Khanh, and G. Zampieri. Propagation of holomorphic extendibility and non-hypoellipticity of the $\bar{\partial}$ -Neumann problem in an exponentially degenerate boundary. *Adv. Math.*, 230(4-6):1972–1978, 2012.
- [BM95] Denis R. Bell and Salah Eldin A. Mohammed. An extension of Hörmander’s theorem for infinitely degenerate second-order operators. *Duke Math. J.*, 78(3):453–475, 1995.
- [Boa87] H. Boas. The Szegő projection: Sobolev estimates in regular domains. *Trans. Amer. Math. Soc.*, 300(1):109–132, 1987.
- [BPZ15] L. Baracco, S. Pinton, and G. Zampieri. Hypoellipticity of the Kohn-Laplacian $\square_b$ and of the $\bar{\partial}$ -Neumann problem by means of subelliptic multipliers. *Math. Ann.*, 362(3-4):887–901, 2015. <http://dx.doi.org/10.1007/s00208-014-1144-1>.
- [Cat83] D. Catlin. Necessary conditions for subellipticity of the $\bar{\partial}$ -Neumann problem. *Ann. of Math.*, 117:147–171, 1983.

- [Cat84] D. Catlin. Global regularity of the $\bar{\partial}$ -Neumann problem. In *Complex analysis of several variables (Madison, Wis., 1982)*, Proc. Sympos. Pure Math., 41, pages 39–49. Amer. Math. Soc., Providence, RI, 1984.
- [Cat87] D. Catlin. Subelliptic estimates for the $\bar{\partial}$ -Neumann problem on pseudoconvex domains. *Ann. of Math.*, 126:131–191, 1987.
- [Chr88] M. Christ. Regularity properties of the $\bar{\partial}_b$ equation on weakly pseudoconvex CR manifolds of dimension 3. *J. Amer. Math. Soc.*, 1:587–646, 1988.
- [Chr01] Michael Christ. Hypocoercivity in the infinitely degenerate regime. In *Complex analysis and geometry (Columbus, OH, 1999)*, volume 9 of *Ohio State Univ. Math. Res. Inst. Publ.*, pages 59–84. de Gruyter, Berlin, 2001.
- [Chr02] M. Christ. Hypocoercivity of the Kohn Laplacian for three-dimensional tubular Cauchy-Riemann structures. *J. Inst. Math. Jussieu*, 1(2):279–291, 2002.
- [CNS92] D.-C. Chang, A. Nagel, and E.M. Stein. Estimates for the $\bar{\partial}$ -Neumann problem in pseudoconvex domains of finite type in $\mathbb{C}^2$. *Acta Math.*, 169:153–228, 1992.
- [Fed71] V. S. Fedii. On a criterion for hypoellipticity. *Math. Sb.*, 14:15–45, 1971.
- [FK72] G. B. Folland and J. J. Kohn. *The Neumann problem for the Cauchy-Riemann Complex*, volume 75 of *Ann. of Math. Stud.* Princeton University Press, Princeton, New Jersey, 1972.
- [FK88a] C. Fefferman and J.J. Kohn. Estimates of kernels on three-dimensional CR manifolds. *Rev. Mat. Iberoamericana*, 4(3-4):355–405, 1988.
- [FK88b] C. Fefferman and J.J. Kohn. Hölder estimates on domains of complex dimension two and on three dimensional CR manifolds. *Adv. in Math.*, 69:233–303, 1988.
- [FKM90] C. Fefferman, J.J. Kohn, and M. Machedon. Hölder estimates on CR manifolds with a diagonalizable Levi form. *Adv. in Math.*, 84:1–90, 1990.
- [FS74] G.B. Folland and E.M. Stein. Estimates for the $\bar{\partial}_b$ -complex and analysis on the Heisenberg group. *Comm. Pure and Appl. Math.*, 27:429–522, 1974.
- [Him95] A. Alexandrou Himonas. On degenerate elliptic operators of infinite type. *Math. Z.*, 220(3):449–460, 1995.
- [HNW10] J. Halfpap, A. Nagel, and S. Wainger. The Bergman and Szegő kernels near points of infinite type. *Pacific J. Math.*, 246(1):75–128, 2010.
- [Hör67] L. Hörmander. Hypocoercive second order differential equations. *Acta Math.*, 119:147–171, 1967.
- [HPR15] P.S. Harrington, M.M. Peloso, and A.S. Raich. Regularity equivalence of the Szegő projection and the complex Green operator. *Proc. Amer. Math. Soc.*, 143(1):353–367, 2015. arXiv:1305.0188.
- [HR11] P. Harrington and A. Raich. Regularity results for $\bar{\partial}_b$ on CR-manifolds of hypersurface type. *Comm. Partial Differential Equations*, 36(1):134–161, 2011.
- [HR15] Phillip S. Harrington and Andrew S. Raich. Closed range for $\bar{\partial}$ and $\bar{\partial}_b$ on bounded hypersurfaces in Stein manifolds. *Ann. Inst. Fourier (Grenoble)*, 65(4):1711–1754, 2015.
- [Ker72] N. Kerzman. The Bergman kernel function. Differentiability at the boundary. *Math. Ann.*, 195:149–158, 1972.
- [Kha16] Tran Vu Khanh. Equivalence of estimates on a domain and its boundary. *Vietnam J. Math.*, 44(1):29–48, 2016.
- [KN65] J.J. Kohn and L. Nirenberg. Non-coercive boundary value problems. *Comm. Pure Appl. Math.*, 18:443–492, 1965.
- [KN06] J.J. Kohn and A. Nicoara. The $\bar{\partial}_b$ -equation on weakly pseudo-convex CR manifolds of dimension 3. *J. Funct. Anal.*, 230:251–272, 2006.

- [Koe02] K. Koenig. On maximal Sobolev and Hölder estimates for the tangential Cauchy-Riemann operator and boundary Laplacian. *Amer. J. Math.*, 124:129–197, 2002.
- [Koh79] J.J. Kohn. Subellipticity of the $\bar{\partial}$ -Neumann problem on pseudoconvex domains: sufficient conditions. *Acta Math.*, 142:79–122, 1979.
- [Koh86] J.J. Kohn. The range of the tangential Cauchy-Riemann operator. *Duke Math. J.*, 53:525–545, 1986.
- [Koh98] J. J. Kohn. Hypoellipticity of some degenerate subelliptic operators. *J. Funct. Anal.*, 159(1):203–216, 1998.
- [Koh00] J. J. Kohn. Hypoellipticity at points of infinite type. In *Analysis, geometry, number theory: the mathematics of Leon Ehrenpreis (Philadelphia, PA, 1998)*, volume 251 of *Contemp. Math.*, pages 393–398. Amer. Math. Soc., Providence, RI, 2000.
- [Koh02] J.J. Kohn. Superlogarithmic estimates on pseudoconvex domains and CR manifolds. *Ann. of Math.*, 156:213–248, 2002.
- [KPZ12] Tran Vu Khanh, Stefano Pinton, and Giuseppe Zampieri. Compactness estimates for $\square_b$ on a CR manifold. *Proc. Amer. Math. Soc.*, 140(9):3229–3236, 2012.
- [KR] Tran Vu Khanh and Andrew Raich. The Kohn-Laplace equation on abstract CR manifolds: global regularity. *submitted*.
- [KS87] S. Kusuoka and D. Stroock. Applications of the Malliavin calculus. III. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 34(2):391–442, 1987.
- [KZ10] T. Khanh and G. Zampieri. Regularity of the $\bar{\partial}$ -Neumann problem at point of infinite type. *J. Funct. Anal.*, 259(11):2760–2775, 2010.
- [KZ12] T. Khanh and G. Zampieri. Necessary geometric and analytic conditions for general estimates in the $\bar{\partial}$ -Neumann problem. *Invent. Math.*, 128(3):729–750, 2012.
- [Mac88] M. Machedon. Estimates for the parametrix of the Kohn Laplacian on certain domains. *Invent. Math.*, 91:339–364, 1988.
- [McN92] J.D. McNeal. Convex domains of finite type. *J. Funct. Anal.*, 108:361–373, 1992.
- [Mor87] Yoshinori Morimoto. Hypoellipticity for infinitely degenerate elliptic operators. *Osaka J. Math.*, 24(1):13–35, 1987.
- [MS97] J.D. McNeal and E.M. Stein. The Szegő projection on convex domains. *Math. Z.*, 224:519–553, 1997.
- [Nic06] A. Nicoara. Global regularity for $\bar{\partial}_b$ on weakly pseudoconvex CR manifolds. *Adv. Math.*, 199:356–447, 2006.
- [NRSW89] A. Nagel, J.-P. Rosay, E.M. Stein, and S. Wainger. Estimates for the Bergman and Szegő kernels in $\mathbb{C}^2$. *Ann. of Math.*, 129:113–149, 1989.
- [NS06] A. Nagel and E.M. Stein. The $\bar{\partial}_b$ -complex on decoupled domains in $\mathbb{C}^n$, $n \geq 3$. *Ann. of Math.*, 164:649–713, 2006.
- [PS77] D.H. Phong and E.M. Stein. Estimates for the Bergman and Szegő projections on strongly pseudo-convex domains. *Duke Math. J.*, 44(3):695–704, 1977.
- [Rai10] A. Raich. Compactness of the complex Green operator on CR-manifolds of hypersurface type. *Math. Ann.*, 348(1):81–117, 2010.
- [RS76] Linda Preiss Rothschild and E. M. Stein. Hypoelliptic differential operators and nilpotent groups. *Acta Math.*, 137(3-4):247–320, 1976.
- [RS08] A. Raich and E. Straube. Compactness of the complex Green operator. *Math. Res. Lett.*, 15(4):761–778, 2008.
- [Sha85] M.-C. Shaw. L^2 -estimates and existence theorems for the tangential Cauchy-Riemann complex. *Invent. Math.*, 82:133–150, 1985.

- [Str10] E. Straube. *Lectures on the $\mathcal{L}^2$ -Sobolev Theory of the $\bar{\partial}$ -Neumann Problem*. ESI Lectures in Mathematics and Physics. European Mathematical Society (EMS), Zürich, 2010.
- [Str12] Emil J. Straube. The complex Green operator on CR-submanifolds of $\mathbb{C}^n$ of hypersurface type: compactness. *Trans. Amer. Math. Soc.*, 364(8):4107–4125, 2012.
- [SZ15] Emil J. Straube and Yunus E. Zeytuncu. Sobolev estimates for the complex Green operator on CR submanifolds of hypersurface type. *Invent. Math.*, 201(3):1073–1095, 2015.

TRAN VU KHANH

SCHOOL OF MATHEMATICS AND APPLIED STATISTICS, UNIVERSITY OF WOLLONGONG, NSW, AUSTRALIA, 2522

E-mail address: tkhanh@uow.edu.au